

TP53I3 rabbit monoclonal antibody

Catalog # H00009540-K Size 100 ug x up to 3

Specification	
Product Description	Rabbit monoclonal antibody raised against a human TP53l3 peptide using ARM Technology.
lmmunogen	A synthetic peptide of human TP53l3 is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
Host	Rabbit
Library Construction	Non-fusion antibody library from rabbit spleen (ARM Technology).
Expression	Overexpression vector and transfection into 293H cell line.
Reactivity	Human
Purification	Protein A
Isotype	lgG
Quality Control Testing	Antibody reactive against human TP53l3 peptide by ELISA and mammalian transfected lysate by W estern Blot.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Deliverable	Up to three rabbit lgG clones of 100 ug each will be delivered to customer.
Note	 Customer may provide cell or tissue lysate for antibody screening. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering in cluding F(ab)₂, lgG, scFv and different Fc and non-Fc conjugates per customer request.

Applications

Western Blot (Transfected lysate)

Protocol Download



ELISA

Gene Info — TP53I3	
Entrez GenelD	9540
GeneBank Accession#	<u>TP53l3</u>
Gene Name	TP53l3
Gene Alias	PIG3
Gene Description	tumor protein p53 inducible protein 3
Omim ID	605171
Gene Ontology	<u>Hyperlink</u>
Gene Summary	The protein encoded by this gene is similar to oxidoreductases, which are enzymes involved in ce llular responses to oxidative stresses and irradiation. This gene is induced by the tumor suppress or p53 and is thought to be involved in p53-mediated cell death. It contains a p53 consensus binding site in its promoter region and a downstream pentanucleotide microsatellite sequence. P53 has been shown to transcriptionally activate this gene by interacting with the downstream pentanucle otide microsatellite sequence. The microsatellite is polymorphic, with a varying number of pentanu cleotide repeats directly correlated with the extent of transcriptional activation by p53. It has been suggested that the microsatellite polymorphism may be associated with differential susceptibility to cancer. At least two transcript variants encoding the same protein have been found for this gene. [provided by RefSeq
Other Designations	OTTHUMP00000116037 p53-induced gene 3 protein quinone oxidoreductase homolog

Pathway

p53 signaling pathway

Disease

- Adenocarcinoma
- Breast cancer
- Breast Neoplasms
- Esophageal Neoplasms



- Genetic Predisposition to Disease
- Hematologic Diseases
- Hodgkin Disease
- <u>Lung Neoplasms</u>
- Lymphoma
- Lymphoproliferative Disorders
- Neoplasms
- Occupational Diseases
- Ovarian cancer
- Ovarian Neoplasms
- Pulmonary Disease
- Urinary Bladder Neoplasms
- Waldenstrom Macroglobulinemia
- Werner syndrome