


RNF10 293T Cell Transient Overexpression Lysate(Denatured)


Catalog # H00009921-T02 Size 100 uL

Applications



SDS-PAGE Gel

RNF10 transfected lysate.



Western Blot

Lane 1: RNF10 transfected lysate (89.32 KDa)

Lane 2: Non-transfected lysate.

| Specification | |
|----------------------------------|------------------------|
| Transfected Cell Line | 293T |
| Plasmid | pCMV-RNF10 full-length |
| Host | Human |
| Theoretical MW (kDa) | 89.32 |
| Interspecies Antigen Sequence | Mouse (92); Rat (92) |



Product Information

| Quality Control Testing | Transient overexpression cell lysate was tested with Anti-RNF10 antibody (H00009921-B02) by Wes tern Blots. SDS-PAGE Gel RNF10 transfected lysate. Western Blot Lane 1: RNF10 transfected lysate (89.32 KDa) Lane 2: Non-transfected lysate. |
|-------------------------|--|
| Storage Buffer | 1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bro mophenol blue) |
| Storage Instruction | Store at -80°C. Aliquot to avoid repeated freezing and thawing. |

Applications

Western Blot

| Gene Info — RNF10 | |
|---------------------|---|
| Entrez GenelD | 9921 |
| GeneBank Accession# | NM_014868.3 |
| Protein Accession# | NP_055683.3 |
| Gene Name | RNF10 |
| Gene Alias | KIAA0262, MGC126758, MGC126764, RIE2 |
| Gene Description | ring finger protein 10 |
| Gene Ontology | <u>Hyperlink</u> |
| Gene Summary | The protein encoded by this gene contains a ring finger motif, which is known to be involved in protein-protein interactions. The specific function of this protein has not yet been determined. EST data suggests the existence of multiple alternatively spliced transcript variants, however, their full length nature is not known. [provided by RefSeq |
| Other Designations | - |