## FGF19 rabbit monoclonal antibody

Catalog # H00009965-K

ocification

Size 100 ug x up to 3

Specification	
Product Description	Rabbit monoclonal antibody raised against a human FGF19 peptide using ARM Technology.
Immunogen	A synthetic peptide of human FGF19 is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
Host	Rabbit
Library Construction	Non-fusion antibody library from rabbit spleen ( <u>ARM Technology</u> ).
Expression	Overexpression vector and transfection into 293H cell line.
Reactivity	Human
Purification	Protein A
lsotype	lgG
Quality Control Testing	Antibody reactive against human FGF19 peptide by ELISA and mammalian transfected lysate by W estern Blot.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Deliverable	Up to three rabbit IgG clones of 100 ug each will be delivered to customer.
Note	<ol> <li>Customer may provide cell or tissue lysate for antibody screening.</li> <li>Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering in cluding F(ab)<sub>2</sub>, lgG, scFv and different Fc and non-Fc conjugates per customer request.</li> </ol>

## Applications

• Western Blot (Transfected lysate)

Protocol Download

• ELISA

## Gene Info — FGF19

Entrez GenelD	<u>9965</u>
GeneBank Accession#	<u>FGF19</u>
Gene Name	FGF19
Gene Alias	-
Gene Description	fibroblast growth factor 19
Omim ID	<u>603891</u>
Gene Ontology	Hyperlink
Gene Summary	The protein encoded by this gene is a member of the fibroblast growth factor (FGF) family. FGF f amily members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes including embryonic development cell growth, morphogenesis, tissue rep air, tumor growth and invasion. This growth factor is a high affinity, heparin dependent ligand for F GFR4. Expression of this gene was detected only in fetal but not adult brain tissue. Synergistic int eraction of the chick homolog and Wnt-8c has been shown to be required for initiation of inner ear development. [provided by RefSeq
Other Designations	-

## Pathway

- MAPK signaling pathway
- <u>Melanoma</u>
- Pathways in cancer
- Regulation of actin cytoskeleton