

## TANK rabbit monoclonal antibody

Catalog # H00010010-K Size 100 ug x up to 3

Considiration	
Specification	
Product Description	Rabbit monoclonal antibody raised against a human TANK peptide using ARM Technology.
Immunogen	A synthetic peptide of human TANK is used for rabbit immunization.  Customer or Abnova will decide on the preferred peptide sequence.
Host	Rabbit
Library Construction	Non-fusion antibody library from rabbit spleen ( <u>ARM Technology</u> ).
Expression	Overexpression vector and transfection into 293H cell line.
Reactivity	Human
Purification	Protein A
Isotype	lgG
Quality Control Testing	Antibody reactive against human TANK peptide by ELISA and mammalian transfected lysate by We stern Blot.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Deliverable	Up to three rabbit lgG clones of 100 ug each will be delivered to customer.
Note	<ol> <li>Customer may provide cell or tissue lysate for antibody screening.</li> <li>Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering in cluding F(ab)<sub>2</sub>, lgG, scFv and different Fc and non-Fc conjugates per customer request.</li> </ol>

## **Applications**

Western Blot (Transfected lysate)

Protocol Download



ELISA

Gene Info — TANK	
Entrez GenelD	10010
GeneBank Accession#	TANK
Gene Name	TANK
Gene Alias	I-TRAF, TRAF2
Gene Description	TRAF family member-associated NFKB activator
Omim ID	603893
Gene Ontology	<u>Hyperlink</u>
Gene Summary	The TRAF (tumor necrosis factor receptor-associated factor) family of proteins associate with an d transduce signals from members of the tumor necrosis factor receptor superfamily. The protein encoded by this gene is found in the cytoplasm and can bind to TRAF1, TRAF2, or TRAF3, there by inhibiting TRAF function by sequestering the TRAFs in a latent state in the cytoplasm. For exa mple, the protein encoded by this gene can block TRAF2 binding to LMP1, the Epstein-Barr virus transforming protein, and inhibit LMP1-mediated NF-kappa-B activation. Two transcript variants e ncoding different isoforms have been found for this gene. [provided by RefSeq
Other Designations	TRAF family member-associated NF-KAPPA-B activator TRAF interacting protein TANK

## Disease

- Arthritis
- Cardiovascular Diseases
- Diabetes Mellitus
- Edema
- Genetic Predisposition to Disease
- Hematologic Diseases
- Hepatitis C
- Hodgkin Disease



- Lymphoproliferative Disorders
- Multiple Myeloma
- Occupational Diseases
- Osteoporosis
- Waldenstrom Macroglobulinemia
- Werner syndrome