


RNF41 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # H00010193-T01


Size 100 uL

Applications



SDS-PAGE Gel

RNF41 transfected lysate.



Western Blot

Lane 1: RNF41 transfected lysate (34.98 KDa)

Lane 2: Non-transfected lysate.

Specification

Transfected Cell Line 293T

Plasmid pCMV-RNF41 full-length

Host Human

Theoretical MW (kDa) 34.98

Quality Control Testing Transient overexpression cell lysate was tested with Anti-RNF41 antibody ([H00010193-B01](#)) by Western Blots.
 SDS-PAGE Gel
 RNF41 transfected lysate.
 Western Blot
 Lane 1: RNF41 transfected lysate (34.98 KDa)
 Lane 2: Non-transfected lysate.

Storage Buffer

1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction

Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot

Gene Info — RNF41

Entrez GeneID[10193](#)**GeneBank Accession#**[NM_005785.2](#)**Protein Accession#**[NP_005776.1](#)**Gene Name**

RNF41

Gene Alias

MGC45228, NRDP1, SBB103

Gene Description

ring finger protein 41

Gene Ontology[Hyperlink](#)**Gene Summary**

The protein encoded by this gene contains a RING finger, a motif present in a variety of functionally distinct proteins and known to be involved in protein-protein and protein-DNA interactions. The specific function of this protein has not yet been determined. Three alternatively spliced transcript variants encoding two distinct isoforms have been reported. [provided by RefSeq]

Other Designations

neuregulin receptor degradation protein-1

Pathway

- [Endocytosis](#)

Disease

- [Genetic Predisposition to Disease](#)
- [Parkinson Disease](#)