


MaxPah®

KLRG1 purified MaxPab rabbit polyclonal antibody (D01P)

Catalog # H00010219-D01P Size 100 ug

Applications



Western Blot (Transfected lysate)

Western Blot analysis of KLRG1 expression in transfected 293T cell line (<u>H00010219-T02</u>) by KLRG1 MaxPab polyclonal antibody.

Lane 1: KLRG1 transfected lysate(21.20 KDa).

Lane 2: Non-transfected lysate.

Specification	
Product Description	Rabbit polyclonal antibody raised against a full-length human KLRG1 protein.
Immunogen	KLRG1 (NP_005801.3, 1 a.a. ~ 189 a.a) full-length human protein.
Sequence	MTDSVIYSMLELPTATQAQNDYGPQQKSSSSRPSCSCLVAIALGLLTAVLLSVLLYQWILCQGSNY STCASCPSCPDRWMKYGNHCYYFSVEEKDWNSSLEFCLARDSHLLVITDNQEMSLLQVFLSEAF CWIGLRNNSGWRWEDGSPLNFSRISSNSFVQTCGAINKNGLQASSCEVPLHWVCKKVRL
Host	Rabbit
Reactivity	Human
Interspecies Antigen Sequence	Mouse (56); Rat (54)
Quality Control Testing	Antibody reactive against mammalian transfected lysate.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.



Applications

Western Blot (Transfected lysate)

Western Blot analysis of KLRG1 expression in transfected 293T cell line ($\underline{\text{H00010219-T02}}$) by KLRG1 MaxPab polyclonal antibody.

Lane 1: KLRG1 transfected lysate(21.20 KDa).

Lane 2: Non-transfected lysate.

Protocol Download

Gene Info — KLRG1	
Entrez GenelD	10219
GeneBank Accession#	NM_005810
Protein Accession#	NP_005801.3
Gene Name	KLRG1
Gene Alias	2F1, CLEC15A, MAFA, MAFA-2F1, MAFA-L, MAFA-LIKE, MGC13600
Gene Description	killer cell lectin-like receptor subfamily G, member 1
Omim ID	604874
Gene Ontology	<u>Hyperlink</u>
Gene Summary	Natural killer (NK) cells are lymphocytes that can mediate lysis of certain tumor cells and virus-infe cted cells without previous activation. They can also regulate specific humoral and cell-mediated i mmunity. The protein encoded by this gene belongs to the killer cell lectin-like receptor (KLR) family, which is a group of transmembrane proteins preferentially expressed in NK cells. Studies in micce suggested that the expression of this gene may be regulated by MHC class I molecules. Altern atively spliced transcript variants have been reported, but their full-length natures have not yet been determined. [provided by RefSeq
Other Designations	C-type lectin domain family 15, member A mast cell function-associated antigen (ITIM-containing)

Disease

Tobacco Use Disorder