


MaxPab®

BAIAP2 purified MaxPab rabbit polyclonal antibody (D01P)

Catalog # H00010458-D01P

Size 100 ug

Applications



Western Blot (Transfected lysate)

Western Blot analysis of BAIAP2 expression in transfected 293T cell line ([H00010458-T02](#)) by BAIAP2 MaxPab polyclonal antibody.

Lane 1: BAIAP2 transfected lysate(60.90 KDa).

Lane 2: Non-transfected lysate.

Specification

Product Description	Rabbit polyclonal antibody raised against a full-length human BAIAP2 protein.
Immunogen	BAIAP2 (NP_059345.1, 1 a.a. ~ 552 a.a) full-length human protein.
Sequence	MSLSRSEEMHRLTENVYKTIMEQFNPSLRNFIAMGKNYEKALAGVTYAAKGYFDALVKMGELASE SQGSKELGDVLFQMAEVHRQIQNQLEEMLKSFHNELLTQLEQKVELDSRYLSAALKKYQTEQRS KGDALDKCQAEKKLRKKSQGSKNPQKYSDKELQYDAISNKQGELENVSDGYKTALTEERRRF CFLVEKQCAVAKNSAAYHSKGKELLAQKLPLWQQACADPSKIPERAVQLMQQVASNGATLPSA LSASKSNLVISDPIPGAKPLPVPELAPFVGRMSAQESTPIMNGVTGPDGEDYSPWADRKAQAP KSLSPQSQSKLSDSYSNTLPVRKSVTPKNSYATTENKTLPRSSSMAAGLERNGMRVKAIFSHA AGDNSTLLSFKEGDLITLLVPEARDGWHYGESEKTKMRGWFPFSYTRVLDSDGSDRLHMSLQQG KSSSTGNLLDKDDLAIPPPDYGAASRAFPQAQTASGFKQRPYSVAVPAFSQGLDDYGARMSRNP FAHVQLKPTVTNDRCDLSAQGPEGREHGDGSARTLAGR
Host	Rabbit
Reactivity	Human
Interspecies Antigen Sequence	Mouse (95); Rat (95)
Quality Control Testing	Antibody reactive against mammalian transfected lysate.

Storage Buffer

In 1x PBS, pH 7.4

Storage Instruction

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Transfected lysate)

Western Blot analysis of BAIAP2 expression in transfected 293T cell line ([H00010458-T02](#)) by BAIAP2 MaxPab polyclonal antibody.

Lane 1: BAIAP2 transfected lysate(60.90 KDa).

Lane 2: Non-transfected lysate.

[Protocol Download](#)

Gene Info — BAIAP2

Entrez GeneID[10458](#)**GeneBank Accession#**[NM_017451](#)**Protein Accession#**[NP_059345.1](#)**Gene Name**

BAIAP2

Gene Alias

BAP2, IRSP53

Gene Description

BAI1-associated protein 2

Omim ID[605475](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

The protein encoded by this gene has been identified as a brain-specific angiogenesis inhibitor (BAI1)-binding protein. This adaptor protein links membrane bound G-proteins to cytoplasmic effector proteins. This protein functions as an insulin receptor tyrosine kinase substrate and suggests a role for insulin in the central nervous system. It also associates with a downstream effector of Rho small G proteins, which is associated with the formation of stress fibers and cytokinesis. This protein is involved in lamellipodia and filopodia formation in motile cells and may affect neuronal growth-cone guidance. This protein has also been identified as interacting with the dentatorubral-pallidolysian atrophy gene, which is associated with an autosomal dominant neurodegenerative disease. Alternative splicing results in multiple transcript variants encoding distinct isoforms

Other Designations

insulin receptor substrate p53

Pathway

- [Adherens junction](#)
- [Regulation of actin cytoskeleton](#)

Disease

- [Attention Deficit Disorder with Hyperactivity](#)
- [Functional Laterality](#)
- [Genetic Predisposition to Disease](#)