


HIS1 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # H00010614-T01


Size 100 uL

Applications



SDS-PAGE Gel

HEXIM1 transfected lysate



Western Blot

Lane 1: HEXIM1 transfected lysate (39.6 KDa).

Lane 2: Non-transfected lysate.

Specification

Transfected Cell Line	293T
Plasmid	pCMV-HIS1 full-length
Host	Human
Theoretical MW (kDa)	39.6
Interspecies Antigen Sequence	Mouse (85); Rat (84)

Quality Control Testing

Transient overexpression cell lysate was tested with Anti-HIS 1 antibody ([H00010614-B01](#)) by Western Blots.

SDS-PAGE Gel

HEXIM1 transfected lysate

Western Blot

Lane 1: HEXIM1 transfected lysate (39.6 KDa).

Lane 2: Non-transfected lysate.

Storage Buffer

1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction

Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot

Gene Info — HEXIM1

Entrez GeneID[10614](#)**GeneBank Accession#**[BC006460](#)**Protein Accession#**[AAH06460](#)**Gene Name**

HEXIM1

Gene Alias

CLP1, EDG1, FLJ13562, HIS1, MAQ1

Gene Description

hexamethylene bis-acetamide inducible 1

Omim ID[607328](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

Expression of this gene is induced by hexamethylene-bis-acetamide in vascular smooth muscle cells. This gene has no introns. [provided by RefSeq]

Other Designations

HMBA-inducible|hexamethylene bisacetamide-inducible protein|hexamethylene-bis-acetamide-inducible transcript 1|hexamthylene bis-acetamide inducible 1|menage a quatre 1