

## MGEA5 rabbit monoclonal antibody

Catalog # H00010724-K Size 100 ug x up to 3

| Specification           |   |
|-------------------------|---|
| Product Description     | Rabbit monoclonal antibody raised against a human MGEA5 peptide using ARM Technology.   |
| Immunogen               | A synthetic peptide of human MGEA5 is used for rabbit immunization.  Customer or Abnova will decide on the preferred peptide sequence.  |
| Host                    | Rabbit  |
| Library Construction    | Non-fusion antibody library from rabbit spleen (ARM Technology).  |
| Expression              | Overexpression vector and transfection into 293H cell line.   |
| Reactivity              | Human   |
| Purification            | Protein A   |
| Isotype                 | lgG   |
| Quality Control Testing | Antibody reactive against human MGEA5 peptide by ELISA and mammalian transfected lysate by W estern Blot.   |
| Storage Buffer          | In 1x PBS, pH 7.4   |
| Storage Instruction     | Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.  |
| Deliverable             | Up to three rabbit lgG clones of 100 ug each will be delivered to customer.   |
| Note                    | <ol> <li>Customer may provide cell or tissue lysate for antibody screening.</li> <li>Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering in cluding F(ab)<sub>2</sub>, lgG, scFv and different Fc and non-Fc conjugates per customer request.</li> </ol> |

## **Applications**

Western Blot (Transfected lysate)

Protocol Download



ELISA

| Gene Info — MGEA5   |   |
|---------------------|---|
| Entrez GenelD       | 10724   |
| GeneBank Accession# | MGEA5   |
| Gene Name           | MGEA5   |
| Gene Alias          | FLJ11229, FLJ23355, KIAA0679, MEA5, NCOAT, OGA  |
| Gene Description    | meningioma expressed antigen 5 (hyaluronidase)  |
| Omim ID             | 604039  |
| Gene Ontology       | Hyperlink   |
| Gene Summary        | The dynamic modification of cytoplasmic and nuclear proteins by O-linked N-acetylglucosamine (O-GlcNAc) addition and removal on serine and threonine residues is catalyzed by OGT (MIM 300 255), which adds O-GlcNAc, and MGEA5, a glycosidase that removes O-GlcNAc modifications (Gao et al., 2001 [PubMed 11148210]).[supplied by OMIM |
| Other Designations  | O-GlcNAcase OTTHUMP00000020345 hyaluronidase in meningioma nuclear cytoplasmic O-Glc NAcase and acetyltransferase   |

## Disease

- Cardiovascular Diseases
- Diabetes Mellitus
- Edema