

LILRB1 rabbit monoclonal antibody

Catalog # H00010859-K

Size 100 ug x up to 3

Specification

Product Description	Rabbit monoclonal antibody raised against a human LILRB1 peptide using ARM Technology.
Immunogen	A synthetic peptide of human LILRB1 is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
Host	Rabbit
Library Construction	Non-fusion antibody library from rabbit spleen (ARM Technology).
Expression	Overexpression vector and transfection into 293H cell line.
Reactivity	Human
Purification	Protein A
Isotype	IgG
Quality Control Testing	Antibody reactive against human LILRB1 peptide by ELISA and mammalian transfected lysate by Western Blot.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Deliverable	Up to three rabbit IgG clones of 100 ug each will be delivered to customer.
Note	1. Customer may provide cell or tissue lysate for antibody screening. 2. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering including F(ab) ₂ , IgG, scFv and different Fc and non-Fc conjugates per customer request.

Applications

- Western Blot (Transfected lysate)

[Protocol Download](#)

- ELISA

Gene Info — LILRB1

Entrez GeneID	10859
GeneBank Accession#	LILRB1
Gene Name	LILRB1
Gene Alias	CD85, CD85J, FLJ37515, ILT2, LIR-1, LIR1, MIR-7, MIR7
Gene Description	leukocyte immunoglobulin-like receptor, subfamily B (with TM and ITIM domains), member 1
Omim ID	604811
Gene Ontology	Hyperlink
Gene Summary	<p>This gene is a member of the leukocyte immunoglobulin-like receptor (LIR) family, which is found in a gene cluster at chromosomal region 19q13.4. The encoded protein belongs to the subfamily B class of LIR receptors which contain two or four extracellular immunoglobulin domains, a transmembrane domain, and two to four cytoplasmic immunoreceptor tyrosine-based inhibitory motifs (ITIMs). The receptor is expressed on immune cells where it binds to MHC class I molecules on antigen-presenting cells and transduces a negative signal that inhibits stimulation of an immune response. It is thought to control inflammatory responses and cytotoxicity to help focus the immune response and limit autoreactivity. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]</p>
Other Designations	CD85 antigen lg-like transcript 2 OTTHUMP00000068384 OTTHUMP00000068385 OTTHUMP00000068386 OTTHUMP00000068408 immunoglobulin-like transcript 2 leukocyte lg-like receptor-1 leukocyte immunoglobulin-like receptor 1 leukocyte immunoglobulin-like receptor sub

Disease

- [Arthritis](#)
- [Cardiovascular Diseases](#)
- [Diabetes Mellitus](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)