


MaxPab®

HSF2BP MaxPab mouse polyclonal antibody (B01P)

Catalog # H00011077-B01P

Size 50 ug

Applications




Western Blot (Transfected lysate)

Western Blot analysis of HSF2BP expression in transfected 293T cell line ([H00011077-T01](#)) by HSF2BP MaxPab polyclonal antibody.

Lane 1: HSF2BP transfected lysate(36.74 KDa).

Lane 2: Non-transfected lysate.



Immunofluorescence

Immunofluorescence of purified MaxPab antibody to HSF2BP on Hs 181.Tes cell. [antibody concentration 10 ug/ml]

Specification

Product Description

Mouse polyclonal antibody raised against a full-length human HSF2BP protein.

Immunogen

HSF2BP (NP_008962.1, 1 a.a. ~ 334 a.a) full-length human protein.

Sequence

MGEAGAAEEACRHMGTKEEFVKVRKKDLERLTTEVMQIRDFLPRILNGEVLESFQKLKMEKNLE
RKEQELEQLKMDCEHFARLETQADNIREKKEKLALRQQLNEAKQQLLQQAIEYCTEMGAACT
LLWGVSSSEEVVKAILGGDKALKFFSITGQTMESFVKSLDGDVQELDSDESQFVFALAGIMTNVA
AIACGREFLVNSSRVLLDTILQLLGDLPKPGQCTKLKVLMLMSLYNVSINLKGLKYISESPGFIPLLWW
LLSDPDAEVCLHVLRLVQSVVLEPEVFSKSASEFRSSLPLQRILAMSKSRNPRLQTAAQELLEDL
RTLEHNV

Host

Mouse

Reactivity	Human
Interspecies Antigen Sequence	Mouse (88); Rat (88)
Quality Control Testing	Antibody reactive against mammalian transfected lysate.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Transfected lysate)

Western Blot analysis of HSF2BP expression in transfected 293T cell line ([H00011077-T01](#)) by HSF2BP MaxPab polyclonal antibody.

Lane 1: HSF2BP transfected lysate(36.74 KDa).

Lane 2: Non-transfected lysate.

[Protocol Download](#)

- Immunofluorescence

Immunofluorescence of purified MaxPab antibody to HSF2BP on Hs 181.Tes cell. [antibody concentration 10 ug/ml]

Gene Info — HSF2BP

Entrez GeneID	11077
GeneBank Accession#	NM_007031.1
Protein Accession#	NP_008962.1
Gene Name	HSF2BP
Gene Alias	-
Gene Description	heat shock transcription factor 2 binding protein
Omim ID	604554
Gene Ontology	Hyperlink

Gene Summary

HSF2 binding protein (HSF2BP) associates with HSF2. The interaction occurs between the trimerization domain of HSF2 and the amino terminal hydrophilic region of HSF2BP that comprises two leucine zipper motifs. HSF2BP may therefore be involved in modulating HSF2 activation. [provided by RefSeq]

Other Designations

heat shock factor 2 binding protein

Disease

- [Tobacco Use Disorder](#)