


CHEK2 (Human) Matched Antibody Pair

Catalog # H00011200-AP11

Size 1 Set

Applications



Sandwich ELISA detection sensitivity ranging from 0.3 ng/ml to 100 ng/ml.

Specification

Product Description	This antibody pair set comes with a matched antibody pair to detect and quantify the protein level of human CHEK2.
Reactivity	Human
Interspecies Antigen Sequence	Mouse (83); Rat (83)
Quality Control Testing	Standard curve using recombinant protein (H00011200-P01) as an analyte. Sandwich ELISA detection sensitivity ranging from 0.3 ng/ml to 100 ng/ml.
Supplied Product	Antibody pair set content: 1. Capture antibody: rabbit MaxPab® affinity purified polyclonal anti-CHEK2 (100 ug) 2. Detection antibody: mouse monoclonal anti-CHEK2, IgG2a Kappa (20 ug) *Reagents are sufficient for at least 1-2 x 96 well plates using recommended protocols.
Storage Instruction	Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -20°C storage immediately after use.

Applications

- ELISA Pair (Recombinant protein)

[Protocol Download](#)

Gene Info — CHEK2

Entrez GeneID [11200](#)

Gene Name CHEK2

Gene Alias CDS1, CHK2, HuCds1, LFS2, PP1425, RAD53

Gene Description CHK2 checkpoint homolog (S. pombe)

Omim ID [114480](#) [176807](#) [259500](#) [604373](#) [609265](#)

Gene Ontology [Hyperlink](#)

Gene Summary

In response to DNA damage and replication blocks, cell cycle progression is halted through the control of critical cell cycle regulators. The protein encoded by this gene is a cell cycle checkpoint regulator and putative tumor suppressor. It contains a forkhead-associated protein interaction domain essential for activation in response to DNA damage and is rapidly phosphorylated in response to replication blocks and DNA damage. When activated, the encoded protein is known to inhibit CDC25C phosphatase, preventing entry into mitosis, and has been shown to stabilize the tumor suppressor protein p53, leading to cell cycle arrest in G1. In addition, this protein interacts with and phosphorylates BRCA1, allowing BRCA1 to restore survival after DNA damage. Mutations in this gene have been linked with Li-Fraumeni syndrome, a highly penetrant familial cancer phenotype usually associated with inherited mutations in TP53. Also, mutations in this gene are thought to confer a predisposition to sarcomas, breast cancer, and brain tumors. This nuclear protein is a member of the CDS1 subfamily of serine/threonine protein kinases. Three transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]

Other Designations OTTHUMP00000028871|checkpoint-like protein CHK2|protein kinase CHK2|serine/threonine-protein kinase CHK2

Pathway

- [Cell cycle](#)
- [p53 signaling pathway](#)

Disease

- [Adenocarcinoma](#)

- [Adenoma](#)
- [Ataxia telangiectasia](#)
- [Atherosclerosis](#)
- [Barrett Esophagus](#)
- [Brain Neoplasms](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Carcinoma](#)
- [Cell Transformation](#)
- [Colon cancer](#)
- [Colonic Neoplasms](#)
- [Colorectal Neoplasms](#)
- [DNA Damage](#)
- [Endometrial Neoplasms](#)
- [Esophageal Neoplasms](#)
- [Genetic Predisposition to Disease](#)
- [Glioma](#)
- [Head and Neck Neoplasms](#)
- [Hypertension](#)
- [Kidney Diseases](#)
- [Kidney Failure](#)
- [Kidney Neoplasms](#)
- [Laryngeal Neoplasms](#)
- [Leukemia](#)
- [Lung Neoplasms](#)

- [Malignant melanoma](#)
- [Melanoma](#)
- [Meningeal Neoplasms](#)
- [Meningioma](#)
- [Mouth Neoplasms](#)
- [Neoplasm Invasiveness](#)
- [Neoplasm Metastasis](#)
- [Neoplasm Recurrence](#)
- [Neoplasms](#)
- [Neoplastic Syndromes](#)
- [Neuroma](#)
- [Ovarian cancer](#)
- [Ovarian Neoplasms](#)
- [Pancreatic cancer](#)
- [Pancreatic Neoplasms](#)
- [Pharyngeal Neoplasms](#)
- [Prostate cancer](#)
- [Prostatic Neoplasms](#)
- [Pulmonary Disease](#)
- [Skin Neoplasms](#)
- [Urinary Bladder Neoplasms](#)
- [Werner syndrome](#)