

AAK1 rabbit monoclonal antibody

Catalog # H00022848-K Size 100 ug x up to 3

Specification	
Specification	
Product Description	Rabbit monoclonal antibody raised against a human AAK1 peptide using ARM Technology.
Immunogen	A synthetic peptide of human AAK1 is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
Host	Rabbit
Library Construction	Non-fusion antibody library from rabbit spleen (<u>ARM Technology</u>).
Expression	Overexpression vector and transfection into 293H cell line.
Reactivity	Human
Purification	Protein A
Isotype	lgG
Quality Control Testing	Antibody reactive against human AAK1 peptide by ELISA and mammalian transfected lysate by We stern Blot.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Deliverable	Up to three rabbit lgG clones of 100 ug each will be delivered to customer.
Note	 Customer may provide cell or tissue lysate for antibody screening. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering in cluding F(ab)₂, lgG, scFv and different Fc and non-Fc conjugates per customer request.

Applications

Western Blot (Transfected lysate)

Protocol Download



ELISA

Gene Info — AAK1	
Entrez GenelD	<u>22848</u>
GeneBank Accession#	AAK1
Gene Name	AAK1
Gene Alias	DKFZp686F03202, DKFZp686K16132, FLJ23712, FLJ25931, FLJ31060, FLJ42882, FLJ4525 2, KIAA1048, MGC138170, MGC164568, MGC164570
Gene Description	AP2 associated kinase 1
Gene Ontology	<u>Hyperlink</u>
Gene Summary	Adaptor-related protein complex 2 (AP-2 complexes) functions during receptor-mediated endocyt osis to trigger clathrin assembly, interact with membrane-bound receptors, and recruit encodytic a ccessory factors. This gene encodes a member of the SNF1 subfamily of Ser/Thr protein kinases. The protein interacts with and phosphorylates a subunit of the AP-2 complex, which promotes bin ding of AP-2 to sorting signals found in membrane-bound receptors and subsequent receptor end ocytosis. Its kinase activity is stimulated by clathrin. Alternatively spliced transcript variants have b een described, but their biological validity has not been determined. [provided by RefSeq
Other Designations	adaptor-associated kinase 1

Disease

- Parkinson Disease
- Tobacco Use Disorder