


FBXO28 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # H00023219-T01


Size 100 uL

Applications



SDS-PAGE Gel

FBXO28 transfected lysate.



Western Blot

Lane 1: FBXO28 transfected lysate (40.59 KDa)

Lane 2: Non-transfected lysate.

Specification

Transfected Cell Line	293T
Plasmid	pCMV-FBXO28 full-length
Host	Human
Theoretical MW (kDa)	40.59
Interspecies Antigen Sequence	Mouse (94); Rat (95)

Quality Control Testing

Transient overexpression cell lysate was tested with Anti-FBXO28 antibody ([H00023219-B01](#)) by Western Blots.
SDS-PAGE Gel
FBXO28 transfected lysate.
Western Blot
Lane 1: FBXO28 transfected lysate (40.59 KDa)
Lane 2: Non-transfected lysate.

Storage Buffer

1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction

Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot

Gene Info — FBXO28

Entrez GeneID[23219](#)**GeneBank Accession#**[NM_015176.1](#)**Protein Accession#**[NP_055991.1](#)**Gene Name**

FBXO28

Gene Alias

FLJ10766, Fbx28, KIAA0483

Gene Description

F-box protein 28

Omim ID[609100](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

Members of the F-box protein family, such as FBXO28, are characterized by an approximately 40 -amino acid F-box motif. SCF complexes, formed by SKP1 (MIM 601434), cullin (see CUL1; MIM 603134), and F-box proteins, act as protein-ubiquitin ligases. F-box proteins interact with SKP1 through the F box, and they interact with ubiquitination targets through other protein interaction domains (Jin et al., 2004 [PubMed 15520277]).[supplied by OMIM]

Other Designations

OTTHUMP00000035597