


MaxPab®

SIRT3 purified MaxPab mouse polyclonal antibody (B01P)

Catalog # H00023410-B01P

Size 50 ug

Applications



Western Blot (Transfected lysate)

Western Blot analysis of SIRT3 expression in transfected 293T cell line ([H00023410-T01](#)) by SIRT3 MaxPab polyclonal antibody.

Lane 1: SIRT3 transfected lysate(44 KDa).

Lane 2: Non-transfected lysate.

Specification

Product Description	Mouse polyclonal antibody raised against a full-length human SIRT3 protein.
Immunogen	SIRT3 (AAH01042, 1 a.a. ~ 399 a.a) full-length human protein.
Sequence	MAFWGWRAAAALRLWGRVVERVEAGGGVGPFQACGCRLVLGGRDDVSAGLRGSHGARGEPLDPVRPLQRPPRPEVPRAFRRQPRAAAPSFSSIKGGRRSISFSVGASSVVGSGGSSDKGKLSLQDVAELIRARACQRVVVMVGAGISTPSGIPDFRSPGSGLYSNLQQYDLPYPEAIFELPFFFHNPKPFFTLAKELYPGNYKPNVTHYFLRLLHDKGLLLRLYTQNIIDGLERVSGIPASKLVEAHGTFASATCTVCQRPFGEDIRADVMDRVPRCPVCTGVVKPDMFFGEPLPQRFLHVVDFPMADLLLILGTSLEVEPFASLTEAVRSSVPRLLINRDLVGPLAWHPRSRDVAQLGDVVHGVESLVELLWTEEMRDLVQRETGKLDGPKD
Host	Mouse
Reactivity	Human
Interspecies Antigen Sequence	Mouse (83); Rat (85)
Quality Control Testing	Antibody reactive against mammalian transfected lysate.
Storage Buffer	In 1x PBS, pH 7.4

Storage Instruction

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Transfected lysate)

Western Blot analysis of SIRT3 expression in transfected 293T cell line ([H00023410-T01](#)) by SIRT3 MaxPab polyclonal antibody.

Lane 1: SIRT3 transfected lysate(44 KDa).

Lane 2: Non-transfected lysate.

[Protocol Download](#)

Gene Info — SIRT3

Entrez GeneID [23410](#)

GeneBank Accession# [BC001042](#)

Protein Accession# [AAH01042](#)

Gene Name SIRT3

Gene Alias SIR2L3

Gene Description sirtuin (silent mating type information regulation 2 homolog) 3 (S. cerevisiae)

Omim ID [604481](#)

Gene Ontology [Hyperlink](#)

Gene Summary This gene encodes a member of the sirtuin family of proteins, homologs to the yeast Sir2 protein. Members of the sirtuin family are characterized by a sirtuin core domain and grouped into four classes. The functions of human sirtuins have not yet been determined; however, yeast sirtuin proteins are known to regulate epigenetic gene silencing and suppress recombination of rDNA. Studies suggest that the human sirtuins may function as intracellular regulatory proteins with mono-ADP-ribosyltransferase activity. The protein encoded by this gene is included in class I of the sirtuin family. Two alternatively spliced transcript variants that encode different proteins have been described for this gene. [provided by RefSeq]

Other Designations mitochondrial nicotinamide adenine dinucleotide-dependent deacetylase|silent mating type information regulation 2, S.cerevisiae, homolog 3|sir2-like 3|sirtuin 3|sirtuin type 3

Disease

- [Alzheimer disease](#)
- [Celiac Disease](#)
- [Diabetes Mellitus](#)
- [Genetic Predisposition to Disease](#)
- [Head and Neck Neoplasms](#)
- [Neoplasm Recurrence](#)
- [Neoplasms](#)