


## SIRT3 293T Cell Transient Overexpression Lysate(Denatured)


Catalog # H00023410-T02 Size 100 uL

## **Applications**



#### SDS-PAGE Gel

SIRT3 transfected lysate.



#### Western Blot

Lane 1: SIRT3 transfected lysate (28.6 KDa)

Lane 2: Non-transfected lysate.

| Specification                    |                        |
|----------------------------------|------------------------|
| Transfected Cell Line            | 293T                   |
| Plasmid                          | pCMV-SIRT3 full-length |
| Host                             | Human                  |
| Theoretical MW (kDa)             | 28.6                   |
| Interspecies Antigen<br>Sequence | Mouse (83); Rat (85)   |



### **Product Information**

| Quality Control Testing | Transient overexpression cell lysate was tested with Anti-SIRT3 antibody (H00023410-B02) by West ern Blots.  SDS-PAGE Gel  SIRT3 transfected lysate.  Western Blot  Lane 1: SIRT3 transfected lysate (28.6 KDa)  Lane 2: Non-transfected lysate. |
|-------------------------|--|
| Storage Buffer          | 1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bro mophenol blue)   |
| Storage Instruction     | Store at -80°C. Aliquot to avoid repeated freezing and thawing.  |

# Applications

Western Blot

| Gene Info — SIRT3   |  |
|---------------------|--|
| Entrez GenelD       | 23410  |
| GeneBank Accession# | NM_001017524.1   |
| Protein Accession#  | =  |
| Gene Name           | SIRT3  |
| Gene Alias          | SIR2L3   |
| Gene Description    | sirtuin (silent mating type information regulation 2 homolog) 3 (S. cerevisiae)  |
| Omim ID             | 604481   |
| Gene Ontology       | <u>Hyperlink</u>   |
| Gene Summary        | This gene encodes a member of the sirtuin family of proteins, homologs to the yeast Sir2 protein. Members of the sirtuin family are characterized by a sirtuin core domain and grouped into four cla sses. The functions of human sirtuins have not yet been determined; however, yeast sirtuin protein s are known to regulate epigenetic gene silencing and suppress recombination of rDNA. Studies suggest that the human sirtuins may function as intracellular regulatory proteins with mono-ADP-ri bosyltransferase activity. The protein encoded by this gene is included in class I of the sirtuin family. Two alternatively spliced transcript variants that encode different proteins have been described for this gene. [provided by RefSeq |
| Other Designations  | mitochondrial nicotinamide adenine dinucleotide-dependent deacetylase silent mating type information regulation 2, S.cerevisiae, homolog 3 sir2-like 3 sirtuin 3 sirtuin type 3  |



### Disease

- Alzheimer disease
- Celiac Disease
- Diabetes Mellitus
- Genetic Predisposition to Disease
- Head and Neck Neoplasms
- Neoplasm Recurrence
- Neoplasms