


MKRN2 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # H00023609-T01


Size 100 uL

Applications



SDS-PAGE Gel

MKRN2 transfected lysate.



Western Blot

Lane 1: MKRN2 transfected lysate (45.87 KDa)

Lane 2: Non-transfected lysate.

Specification

Transfected Cell Line	293T
Plasmid	pCMV-MKRN2 full-length
Host	Human
Theoretical MW (kDa)	45.87
Interspecies Antigen Sequence	Mouse (88); Rat (88)

Quality Control Testing

Transient overexpression cell lysate was tested with Anti-MKRN2 antibody ([H00023609-B01](#)) by Western Blots.
SDS-PAGE Gel
MKRN2 transfected lysate.
Western Blot
Lane 1: MKRN2 transfected lysate (45.87 KDa)
Lane 2: Non-transfected lysate.

Storage Buffer

1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction

Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot

Gene Info — MKRN2

Entrez GeneID[23609](#)**GeneBank Accession#**[NM_014160.3](#)**Protein Accession#**[NP_054879.3](#)**Gene Name**

MKRN2

Gene Alias

HSPC070, RNF62

Gene Description

makorin ring finger protein 2

Omim ID[608426](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

Members of the makorin family, including MKRN2, have a characteristic zinc finger composition that suggests that they are ribonucleoproteins (Gray et al., 2001 [PubMed 11597136]).[supplied by OMIM]

Other Designations

-

Disease

- [Disease Progression](#)

- [Disease Susceptibility](#)
- [HIV Infections](#)