


MaxPab®

QPCT purified MaxPab rabbit polyclonal antibody (D01P)

Catalog # H00025797-D01P


Size 100 ug

Applications



Western Blot (Cell lysate)

QPCT MaxPab rabbit polyclonal antibody. Western Blot analysis of QPCT expression in HeLa.



Western Blot (Transfected lysate)

Western Blot analysis of QPCT expression in transfected 293T cell line ([H00025797-T02](#)) by QPCT MaxPab polyclonal antibody.

Lane 1: QPCT transfected lysate(40.90 KDa).

Lane 2: Non-transfected lysate.

Specification

Product Description

Rabbit polyclonal antibody raised against a full-length human QPCT protein.

Immunogen

QPCT (NP_036545.1, 1 a.a. ~ 361 a.a) full-length human protein.

Sequence

MAGGRHRRVVGTLHLLLLVAALPWASRGVSPSASAWPEEKNYHQPAILNSSALRQIAEGTSISEM
WQNDLQPLLIERYPGSPGSYAARQHIMQRIQLQADWVLEIDTFLSQTPYGYRSFSNIISTLNPTAKR
HLVLACHYDSKYFSHWNNRVFVGATDSAVPCAMMLELARALDKLLSLKTVSDSKPDLSLQLIFF
DGEEAFLHWSPQDSL YGSRHLAAKMASTPHPPGARGTSQLHGMDLLVLLDLIGAPNPTFPNFFP
NSARWFERLQAIEHELHELGLLKDHSLEGRYFQNYSYGGVIQDDHIPFLRRGVPVLHLIPSPFPEV
WHTMDDNEENLDESTIDNLNKILQVFVLEYLHL

Host

Rabbit

Reactivity	Human
Interspecies Antigen Sequence	Mouse (82); Rat (69)
Quality Control Testing	Antibody reactive against mammalian transfected lysate.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Cell lysate)

QPCT MaxPab rabbit polyclonal antibody. Western Blot analysis of QPCT expression in HeLa.

[Protocol Download](#)

- Western Blot (Transfected lysate)

Western Blot analysis of QPCT expression in transfected 293T cell line ([H00025797-T02](#)) by QPCT MaxPab polyclonal antibody.

Lane 1: QPCT transfected lysate(40.90 KDa).

Lane 2: Non-transfected lysate.

[Protocol Download](#)

Gene Info — QPCT

Entrez GeneID	25797
GeneBank Accession#	NM_012413
Protein Accession#	NP_036545.1
Gene Name	QPCT
Gene Alias	GCT, QC
Gene Description	glutamyl-peptide cyclotransferase
Omim ID	607065
Gene Ontology	Hyperlink

Gene Summary

This gene encodes human pituitary glutaminy cyclase, which is responsible for the presence of pyroglutamyl residues in many neuroendocrine peptides. The amino acid sequence of this enzyme is 86% identical to that of bovine glutaminy cyclase. [provided by RefSeq]

Other Designations

glutaminy cyclase

Publication Reference

- [Glutaminy Cyclase As a Diagnostic/Prognostic Indicator For Neurodegenerative Diseases.](#)

Hans-Ulrich Demuth, Stephan Schilling, Martin Kleinschmidt, Jens-Ulrich Rahfeld, Astrid Kehlen, Monique Haegele.

United States Patent Application Publication 2010 Feb; [Epub].

Application: WB-Ti, Human, Brain

Disease

- [Genetic Predisposition to Disease](#)
- [Hypertension](#)
- [Osteoporosis](#)