


ATP2C1 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # H00027032-T02


Size 100 uL

Applications



SDS-PAGE Gel

ATP2C1 transfected lysate.



Western Blot

Lane 1: ATP2C1 transfected lysate (100.60 KDa)

Lane 2: Non-transfected lysate.

Specification

Transfected Cell Line 293T

Plasmid pCMV-ATP2C1 full-length

Host Human

Theoretical MW (kDa) 100.6

Quality Control Testing Transient overexpression cell lysate was tested with Anti-ATP2C1 antibody ([H00027032-D01P](#)) by Western Blots.
SDS-PAGE Gel
ATP2C1 transfected lysate.
Western Blot
Lane 1: ATP2C1 transfected lysate (100.60 KDa)
Lane 2: Non-transfected lysate.

| | |
|---------------------|---|
| Storage Buffer | 1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue) |
| Storage Instruction | Store at -80°C. Aliquot to avoid repeated freezing and thawing. |

Applications

- Western Blot

Gene Info — ATP2C1

| | |
|---------------------|--|
| Entrez GeneID | 27032 |
| GeneBank Accession# | NM_014382 |
| Protein Accession# | NP_055197.2 |
| Gene Name | ATP2C1 |
| Gene Alias | ATP2C1A, BCPM, HHD, KIAA1347, PMR1, SPCA1, hSPCA1 |
| Gene Description | ATPase, Ca ⁺⁺ transporting, type 2C, member 1 |
| Omim ID | 169600 604384 |
| Gene Ontology | Hyperlink |
| Gene Summary | The protein encoded by this gene belongs to the family of P-type cation transport ATPases. This magnesium-dependent enzyme catalyzes the hydrolysis of ATP coupled with the transport of the calcium. Defects in this gene cause Hailey-Hailey disease, an autosomal dominant disorder. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq] |
| Other Designations | ATP-dependent Ca(2+) pump ATPase 2C1 ATPase, Ca(2+)-sequestering HUSY-28 calcium-transporting ATPase 2C1 secretory pathway Ca2+/Mn2+ ATPase |