


PYCARD 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # H00029108-T01 Size 100 uL

Applications



SDS-PAGE Gel

PYCARD transfected lysate

Western Blot

Lane 1: PYCARD transfected lysate (16.5 KDa). Lane 2: Non-transfected lysate.

Specification	
Transfected Cell Line	293T
Plasmid	pCMV-PYCARD full-length
Host	Human
Theoretical MW (kDa)	16.5
Interspecies Antigen Sequence	Mouse (71); Rat (69)



Product Information

Quality Control Testing	Transient overexpression cell lysate was tested with Anti-PYCARD antibody (<u>H00029108-B01</u>) by W estern Blots. SDS-PAGE Gel PYCARD transfected lysate
	Western Blot
	Lane 1: PYCARD transfected lysate (16.5 KDa).
	Lane 2: Non-transfected lysate.
Storage Buffer	1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bro mophenol blue)
Storage Instruction	Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

• Western Blot

Gene Info — PYCARD

Entrez GenelD	<u>29108</u>
GeneBank Accession#	<u>BC013569</u>
Protein Accession#	AAH13569
Gene Name	PYCARD
Gene Alias	ASC, CARD5, MGC10332, TMS, TMS-1, TMS1
Gene Description	PYD and CARD domain containing
Omim ID	<u>606838</u>
Gene Ontology	Hyperlink
Gene Summary	Hyperlink This gene encodes an adaptor protein that is composed of two protein-protein interaction domain s: a N-terminal PYRIN-PAAD-DAPIN domain (PYD) and a C-terminal caspase-recruitment domai n (CARD). The PYD and CARD domains are members of the six-helix bundle death domain-fold superfamily that mediates assembly of large signaling complexes in the inflammatory and apoptot ic signaling pathways via the activation of caspase. In normal cells, this protein is localized to the cytoplasm; however, in cells undergoing apoptosis, it forms ball-like aggregates near the nuclear periphery. Two transcript variants encoding different isoforms have been found for this gene. [prov ided by RefSeq



Disease

- Arthritis
- Genetic Predisposition to Disease