

DNAxPAb

Hard-to-Find  
Antibody

# BLNK DNAxPab

Catalog # H00029760-W01P

Size 200 ug

## Specification

|                         |  |
|-------------------------|--|
| Product Description     | Rabbit polyclonal antibody raised against a partial-length human BLNK DNA using DNAx™ Immune technology. |
| Technology              | <a href="#">DNAx™ Immune</a>   |
| Immunogen               | Extracellular membrane domain (ECD) human DNA  |
| Host                    | Rabbit   |
| Reactivity              | Human  |
| Purification            | Protein A  |
| Quality Control Testing | Antibody reactive against mammalian transfected lysate.  |
| Storage Buffer          | In 1x PBS, pH 7.4  |
| Storage Instruction     | Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.                                 |

## Applications

- Western Blot (Transfected lysate)

[Protocol Download](#)

- Immunofluorescence (Transfected cell)
- Flow Cytometry (Transfected cell)

## Gene Info — BLNK

|                     |   |
|---------------------|---|
| Entrez GeneID       | <a href="#">29760</a>   |
| GeneBank Accession# | <a href="#">BC018906.2</a>  |
| Protein Accession#  | <a href="#">AAH18906.1</a>  |
| Gene Name           | BLNK  |
| Gene Alias          | BASH, BLNK-S, LY57, MGC111051, SLP-65, SLP65  |
| Gene Description    | B-cell linker   |
| Omim ID             | <a href="#">604515</a>  |
| Gene Ontology       | <a href="#">Hyperlink</a>   |
| Gene Summary        | This gene encodes a cytoplasmic linker or adaptor protein that plays a critical role in B cell development. This protein bridges B cell receptor-associated kinase activation with downstream signaling pathways, thereby affecting various biological functions. The phosphorylation of five tyrosine residues is necessary for this protein to nucleate distinct signaling effectors following B cell receptor activation. Mutations in this gene cause hypoglobulinemia and absent B cells, a disease in which the pro- to pre-B-cell transition is developmentally blocked. Deficiency in this protein has also been shown in some cases of pre-B acute lymphoblastic leukemia. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq] |
| Other Designations  | B cell linker protein B-cell adapter containing a SH2 domain protein B-cell adapter containing a Src homology 2 domain protein OTTHUMP00000020167 Src homology 2 domain-containing leukocyte protein of 65 kDa  |

## Pathway

- [B cell receptor signaling pathway](#)
- [Primary immunodeficiency](#)

## Disease

- [Alzheimer Disease](#)
- [Genetic Predisposition to Disease](#)
- [Tobacco Use Disorder](#)