


## SNX15 293T Cell Transient Overexpression Lysate(Denatured)


Catalog # H00029907-T02 Size 100 uL

### **Applications**



#### SDS-PAGE Gel

SNX15 transfected lysate.



#### Western Blot

Lane 1: SNX15 transfected lysate (38.3 KDa)

Lane 2: Non-transfected lysate.

Specification	
Transfected Cell Line	293T
Plasmid	pCMV-SNX15 full-length
Host	Human
Theoretical MW (kDa)	37.73
Interspecies Antigen Sequence	Mouse (88); Rat (86)



### **Product Information**

Quality Control Testing	Transient overexpression cell lysate was tested with Anti-SNX15 antibody (H00029907-B02) by Wes tern Blots.  SDS-PAGE Gel  SNX15 transfected lysate.  Western Blot  Lane 1: SNX15 transfected lysate (38.3 KDa)  Lane 2: Non-transfected lysate.
Storage Buffer	1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bro mophenol blue)
Storage Instruction	Store at -80°C. Aliquot to avoid repeated freezing and thawing.

# Applications

Western Blot

Gene Info — SNX15	
Entrez GenelD	<u>29907</u>
GeneBank Accession#	NM_013306
Protein Accession#	NP_037438
Gene Name	SNX15
Gene Alias	HSAF001435
Gene Description	sorting nexin 15
Omim ID	605964
Gene Ontology	<u>Hyperlink</u>
Gene Summary	This gene encodes a member of the sorting nexin family. Members of this family contain a phox (PX) domain, which is a phosphoinositide binding domain, and are involved in intracellular trafficking. Overexpression of this gene results in a decrease in the processing of insulin and hepatocyte growth factor receptors to their mature subunits. This decrease is caused by the mislocalization of furin, the endoprotease responsible for cleavage of insulin and hepatocyte growth factor receptors. This protein is involved in endosomal trafficking from the plasma membrane to recycling endosomes or the trans-Golgi network. This gene encodes two transcript variants encoding distinct isoforms. [provided by RefSeq
Other Designations	OTTHUMP00000035515 clone iota unknown protein