


CD209 (Human) IP-WB Antibody Pair

Catalog # H00030835-PW1

Size 1 Set

Applications



Immunoprecipitation of CD209 transfected lysate using rabbit polyclonal anti-CD209 and Protein A Magnetic Bead ([U0007](#)), and immunoblotted with mouse purified polyclonal anti-CD209.

Specification

Product Description	This IP-WB antibody pair set comes with one antibody for immunoprecipitation and another to detect the precipitated protein in western blot.
Reactivity	Human
Interspecies Antigen Sequence	Mouse (66); Rat (62)
Quality Control Testing	Immunoprecipitation-Western Blot (IP-WB) Immunoprecipitation of CD209 transfected lysate using rabbit polyclonal anti-CD209 and Protein A Magnetic Bead (U0007), and immunoblotted with mouse purified polyclonal anti-CD209.
Supplied Product	Antibody pair set content: 1. Antibody pair for IP: rabbit polyclonal anti-CD209 (300 ul) 2. Antibody pair for WB: mouse purified polyclonal anti-CD209 (50 ug)
Storage Instruction	Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -20°C storage immediately after use.

Applications

- Immunoprecipitation-Western Blot

[Protocol Download](#)

Gene Info — CD209

Entrez GeneID [30835](#)

Gene Name CD209

Gene Alias CDSIGN, CLEC4L, DC-SIGN, DC-SIGN1, MGC129965

Gene Description CD209 molecule

Omim ID [604672](#) [607948](#) [609423](#)

Gene Ontology [Hyperlink](#)

Gene Summary

This gene encodes a transmembrane receptor and is often referred to as DC-SIGN because of its expression on the surface of dendritic cells and macrophages. The encoded protein is involved in the innate immune system and recognizes numerous evolutionarily divergent pathogens ranging from parasites to viruses with a large impact on public health. The protein is organized into three distinct domains: an N-terminal transmembrane domain, a tandem-repeat neck domain and C-type lectin carbohydrate recognition domain. The extracellular region consisting of the C-type lectin and neck domains has a dual function as a pathogen recognition receptor and a cell adhesion receptor by binding carbohydrate ligands on the surface of microbes and endogenous cells. The neck region is important for homo-oligomerization which allows the receptor to bind multivalent ligands with high avidity. Variations in the number of 23 amino acid repeats in the neck domain of this protein are rare but have a significant impact on ligand binding ability. This gene is closely related in terms of both sequence and function to a neighboring gene (GeneID 10332; often referred to as L-SIGN). DC-SIGN and L-SIGN differ in their ligand-binding properties and distribution. Alternative splicing results in multiple variants

Other Designations C-type lectin domain family 4, member L|CD209 antigen|Dendritic cell-specific intracellular adhesion molecules (ICAM)-3 grabbing non-integrin

Disease

- [Acquired Immunodeficiency Syndrome](#)
- [Arthritis](#)
- [Asthma](#)
- [Birth Weight](#)

- [Bronchiolitis](#)
- [Celiac Disease](#)
- [Colitis](#)
- [Crohn Disease](#)
- [Cytomegalovirus Infections](#)
- [Dengue](#)
- [Disease Progression](#)
- [Disease Susceptibility](#)
- [Genetic Predisposition to Disease](#)
- [Glioblastoma](#)
- [Glioma](#)
- [Hemophilia A](#)
- [Hepatitis C](#)
- [HIV Infections](#)
- [HTLV-I Infections](#)
- [Hypersensitivity](#)
- [Infant](#)
- [Leukemia](#)
- [Liver Cirrhosis](#)
- [Liver Diseases](#)
- [Lymphadenitis](#)
- [Meningeal Neoplasms](#)
- [Meningioma](#)
- [Mycobacterium Infections](#)
- [Nasopharyngeal Neoplasms](#)
- [Periodontitis](#)

- [Respiratory Syncytial Virus Infections](#)
- [Severe Acute Respiratory Syndrome](#)
- [Sexually Transmitted Diseases](#)
- [Tuberculosis](#)