TAS2R10 rabbit monoclonal antibody

Catalog # H00050839-K

Size 100 ug x up to 3

| 0 | ! C ! | |
|----|--------------|--|
| Sh | ecification | |
| Uρ | Combation | |
| | | |

| Product Description | Rabbit monoclonal antibody raised against a human TAS2R10 peptide using ARM Technology. |
|--|--|
| Immunogen | A synthetic peptide of human TAS2R10 is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence. |
| Host | Rabbit |
| Library Construction | Non-fusion antibody library from rabbit spleen (ARM Technology). |
| Expression | Overexpression vector and transfection into 293H cell line. |
| Reactivity | Human |
| Purification | Protein A |
| | |
| lsotype | lgG |
| Isotype Quality Control Testing | IgG Antibody reactive against human TAS2R10 peptide by ELISA and mammalian transfected lysate by Western Blot. |
| | Antibody reactive against human TAS2R10 peptide by ELISA and mammalian transfected lysate by |
| Quality Control Testing | Antibody reactive against human TAS2R10 peptide by ELISA and mammalian transfected lysate by Western Blot. |
| Quality Control Testing Storage Buffer | Antibody reactive against human TAS2R10 peptide by ELISA and mammalian transfected lysate by Western Blot. |

Applications

• Western Blot (Transfected lysate)

Protocol Download

• ELISA

Gene Info — TAS2R10

| Entrez GenelD | <u>50839</u> |
|---------------------|--|
| GeneBank Accession# | TAS2R10 |
| Gene Name | TAS2R10 |
| Gene Alias | MGC126811, MGC126813, T2R10, TRB2 |
| Gene Description | taste receptor, type 2, member 10 |
| Omim ID | <u>604791</u> |
| Gene Ontology | Hyperlink |
| Gene Summary | This gene product belongs to the family of candidate taste receptors that are members of the G-pr otein-coupled receptor superfamily. These proteins are specifically expressed in the taste receptor r cells of the tongue and palate epithelia. They are organized in the genome in clusters and are ge netically linked to loci that influence bitter perception in mice and humans. In functional expression studies, they respond to bitter tastants. This gene maps to the taste receptor gene cluster on chro mosome 12p13. [provided by RefSeq |
| Other Designations | taste receptor, family B, member 2 |

Pathway

• Taste transduction