


#### MaxPab®

# EXOSC1 purified MaxPab mouse polyclonal antibody (B01P)

Catalog # H00051013-B01P

Size 50 ug


## Applications



#### Western Blot (Transfected lysate)

Western Blot analysis of EXOSC1 expression in transfected 293T cell line (<u>H00051013-T01</u>) by EXOSC1 MaxPab polyclonal antibody.

Lane 1: EXOSC1 transfected lysate(21.56 KDa). Lane 2: Non-transfected lysate.



#### Immunofluorescence

Immunofluorescence of <u>purified</u> MaxPab antibody to EXOSC1 on HeLa cell. [antibody concentration 10 ug/ml]

Specification	
Product Description	Mouse polyclonal antibody raised against a full-length human EXOSC1 protein.
Immunogen	EXOSC1 (AAH22067, 1 a.a. ~ 195 a.a) full-length human protein.
Sequence	MAPPVRYCIPGERLCNLEEGSPGSGTYTRHGYIFSSLAGCLMKSSENGALPVVSVVRETESQLLP DVGAIVTCKVSSINSRFAKVHILYVGSMPLKNSFRGTIRKEDVRATEKDKVEIYKSFRPGDIVLAKVI SLGDAQSNYLLTTAENELGVVVAHSESGIQMVPISWCEMQCPKTHTKEFRKVARVQPEFLQT
Host	Mouse
Reactivity	Human

# 😵 Abnova

### **Product Information**

Interspecies Antigen Sequence	Mouse (95); Rat (95)
Quality Control Testing	Antibody reactive against mammalian transfected lysate.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

## Applications

• Western Blot (Transfected lysate)

Western Blot analysis of EXOSC1 expression in transfected 293T cell line (H00051013-T01) by EXOSC1 MaxPab polyclonal antibody.

Lane 1: EXOSC1 transfected lysate(21.56 KDa). Lane 2: Non-transfected lysate.

Protocol Download

#### Immunofluorescence

Immunofluorescence of purified MaxPab antibody to EXOSC1 on HeLa cell. [antibody concentration 10 ug/ml]

## Gene Info — EXOSC1

Entrez GenelD	<u>51013</u>
GeneBank Accession#	<u>BC022067</u>
Protein Accession#	<u>AAH22067</u>
Gene Name	EXOSC1
Gene Alias	CGI-108, CSL4, CsI4p, SKI4, Ski4p, hCsI4p, p13
Gene Description	exosome component 1
Omim ID	<u>606493</u>
Gene Ontology	Hyperlink
Gene Summary	This gene encodes a core component of the exosome. The mammalian exosome is required for r apid degradation of AU rich element-containing RNAs but not for poly(A) shortening. The associat ion of this protein with the exosome is mediated by protein-protein interactions with ribosomal RN A-processing protein 42 and ribosomal RNA-processing protein 46. [provided by RefSeq



### **Product Information**

**Other Designations** 

3'-5' exoribonuclease CSL4 homolog|CSL4 exosomal core protein homolog|OTTHUMP0000002 0203|exosomal core protein CSL4|homolog of yeast exosomal core protein CSL4

## Pathway

RNA degradation