

DDX47 rabbit monoclonal antibody

Catalog # H00051202-K Size 100 ug x up to 3

Specification	
Product Description	Rabbit monoclonal antibody raised against a human DDX47 peptide using ARM Technology.
Immunogen	A synthetic peptide of human DDX47 is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
Host	Rabbit
Library Construction	Non-fusion antibody library from rabbit spleen (<u>ARM Technology</u>).
Expression	Overexpression vector and transfection into 293H cell line.
Reactivity	Human
Purification	Protein A
Isotype	lgG
Quality Control Testing	Antibody reactive against human DDX47 peptide by ELISA and mammalian transfected lysate by W estern Blot.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Deliverable	Up to three rabbit lgG clones of 100 ug each will be delivered to customer.
Note	 Customer may provide cell or tissue lysate for antibody screening. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering in cluding F(ab)₂, lgG, scFv and different Fc and non-Fc conjugates per customer request.

Applications

Western Blot (Transfected lysate)

Protocol Download



ELISA

Gene Info — DDX47	
Entrez GenelD	<u>51202</u>
GeneBank Accession#	DDX47
Gene Name	DDX47
Gene Alias	DKFZp564O176, E4-DBP, FLJ30012, HQ0256, MSTP162
Gene Description	DEAD (Asp-Glu-Ala-Asp) box polypeptide 47
Gene Ontology	<u>Hyperlink</u>
Gene Summary	This gene encodes a member of the DEAD box protein family. DEAD box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are putative RNA helicases. They are implicate d in a number of cellular processes involving alteration of RNA secondary structure, such as transl ation initiation, nuclear and mitochondrial splicing, and ribosome and spliceosome assembly. Ba sed on their distribution patterns, some members of this family are believed to be involved in emb ryogenesis, spermatogenesis, and cellular growth and division. The protein encoded by this gene can shuttle between the nucleus and the cytoplasm, and has an RNA-independent ATPase activity . Two alternatively spliced transcript variants encoding distinct isoforms have been found for this g ene. [provided by RefSeq
Other Designations	DEAD box polypeptide 47 E4-DEAD box protein