


NOP16 (Human) IP-WB Antibody Pair

Catalog # H00051491-PW1 Size 1 Set

Applications



Immunoprecipitation of NOP16 transfected lysate using rabbit polyclonal anti-NOP16 and Protein A Magnetic Bead (<u>U0007</u>), and immunoblotted with mouse purified polyclonal anti-NOP16.

| Specification | |
|----------------------------------|---|
| Product Description | This IP-WB antibody pair set comes with one antibody for immunoprecipitation and another to detect the precipitated protein in western blot. |
| Reactivity | Human |
| Interspecies Antigen Sequence | Mouse (93); Rat (93) |
| Quality Control Testing | Immunoprecipitation-Western Blot (IP-WB) Immunoprecipitation of NOP16 transfected lysate using rabbit polyclonal anti-NOP16 and Protein A Magnetic Bead (U0007), and immunoblotted with mouse purified polyclonal anti-NOP16. |
| Supplied Product | Antibody pair set content: 1. Antibody pair for IP: rabbit polyclonal anti-NOP16 (300 ul) 2. Antibody pair for WB: mouse purified polyclonal anti-NOP16 (50 ug) |
| Storage Instruction | Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze tha w cycle. Reagents should be returned to -20°C storage immediately after use. |

Applications



• Immunoprecipitation-Western Blot

Protocol Download

| Gene Info — NOP16 | |
|--------------------|--|
| Entrez GenelD | <u>51491</u> |
| Gene Name | NOP16 |
| Gene Alias | HSPC111, HSPC185 |
| Gene Description | NOP16 nucleolar protein homolog (yeast) |
| Gene Ontology | <u>Hyperlink</u> |
| Gene Summary | NOP16 is transcriptionally regulated by c-Myc (MYC; MIM 190080), upregulated in breast cancer, and overexpression is associated with poor patient survival (Butt et al., 2008).[supplied by OMIM |
| Other Designations | HBV pre-S2 trans-regulated protein 3 NOP16 nucleolar protein homolog nucleolar protein 16 homolog |