

DNAxPAb

Hard-to-Find
Antibody

MED15 DNAxPab

Catalog # H00051586-W01P

Size 200 ug

Specification

Product Description Rabbit polyclonal antibody raised against a full-length human MED15 DNA using DNAx™ Immune technology.

Technology [DNAx™ Immune](#)

Immunogen Full-length human DNA

Sequence

MDVSGQETDWRSTAFRQKLVSQIEDAMRKAGVAHSSKSSKDMESHVFLKAKTRDEYLSLVARLIH
FRDIHNKKSQASVSDPMNALQSLTGGPAAGAAGIGMPPRGPGQSLGGMGSLGAMGQPMSLSG
QPPPGTSGMAPHSMVAVSTATPQTQLQLQQVALQQQQQQQQFQQQQQAALQQQQQQQQQQQ
FQAQQSAMQQQFQAVVQQQQQLQQQQQQQHLIKLHHQNQQQIQQQQQQLQRIQLQLQQQQ
QQQQQQQQQQQALQAQPPIQPPMQQPQPPPSQALPQQLQQMHHTQHHQPPQPQPPVA
QNQPSQLPPQSQTQPLVSQAQALPGQMLYTQPPLKFVRAPMVVQQPPVQPQVQQQTAVQTA
QAAQMVAPGVQVSQSSLPMLSSPSPGQVQTPQSMPPPPQSPQPGQPSSQPNVSSGPA
PSPSSFLPSPSPQSPVPTARTPQNFVSPSPGPLNTPVNPSSVMSPAGSSQAEQQYLDKLG
QLSKYIEPLRRMINKIDKNEDRKKDLSKMSLLDILTDPKRCPLKTLQKCEIALEKLNDAVPTP
PPPPVPPTKQQYLCQPLLDVLANIRSPVFNHSLYRTFVPAMTAIHGPITAPVVCTRKRLEDDE
RQSIPSVLQGEVARLDPKFLVNLDPHSCSNNGTVHLICKLDDKDLPSVPPELSVPADYPAQSPL
WIDRQWQYDANPFLQSVHRCMTSRLQLPDKHVSHTALLNTWAQSVHQACLSAA

Host Rabbit

Reactivity Human

Purification Protein A

Quality Control Testing Antibody reactive against mammalian transfected lysate.

Storage Buffer In 1x PBS, pH 7.4

Storage Instruction Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Transfected lysate)

[Protocol Download](#)

- Immunofluorescence (Transfected cell)

- Flow Cytometry (Transfected cell)

Gene Info — MED15

Entrez GeneID [51586](#)

GeneBank Accession# [BC013985.1](#)

Protein Accession# [AAH13985.1](#)

Gene Name MED15

Gene Alias ARC105, CAG7A, CTG7A, DKFZp686A2214, DKFZp762B1216, FLJ42282, FLJ42935, PCQA P, TIG-1, TIG1, TNRC7

Gene Description mediator complex subunit 15

Omim ID [607372](#)

Gene Ontology [Hyperlink](#)

Gene Summary The protein encoded by this gene is a subunit of the multiprotein complexes PC2 and ARC/DRIP and may function as a transcriptional coactivator in RNA polymerase II transcription. This gene contains stretches of trinucleotide repeats and is located in the chromosome 22 region which is deleted in DiGeorge syndrome. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]

Other Designations CTG repeat protein 7a|PC2 (positive cofactor 2, multiprotein complex) glutamine/Q-rich-associated protein|PC2-glutamine-rich-associated protein|TPA inducible gene-1|TPA inducible protein|activator-recruited cofactor, 105-kD|positive cofactor 2, glutamine/

Disease

- [Schizophrenia](#)