


MED9 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # H00055090-T02


Size 100 uL

Applications



SDS-PAGE Gel

MED9 transfected lysate.



Western Blot

Lane 1: MED9 transfected lysate (16.17 KDa)

Lane 2: Non-transfected lysate.

Specification

Transfected Cell Line	293T
Plasmid	pCMV-MED9 full-length
Host	Human
Theoretical MW (kDa)	16.17
Interspecies Antigen Sequence	Mouse (78); Rat (78)

Quality Control Testing

Transient overexpression cell lysate was tested with Anti-MED9 antibody ([H00055090-B01P](#)) by Western Blots.
SDS-PAGE Gel
MED9 transfected lysate.
Western Blot
Lane 1: MED9 transfected lysate (16.17 KDa)
Lane 2: Non-transfected lysate.

Storage Buffer

1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction

Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot

Gene Info — MED9

Entrez GeneID

[55090](#)

GeneBank Accession#

[NM_018019](#)

Protein Accession#

[NP_060489](#)

Gene Name

MED9

Gene Alias

FLJ10193, MED25, MGC138234

Gene Description

mediator complex subunit 9

Omim ID

[609878](#)

Gene Ontology

[Hyperlink](#)

Gene Summary

The multiprotein Mediator complex is a coactivator required for activation of RNA polymerase II transcription by DNA bound transcription factors. The protein encoded by this gene is thought to be a subunit of the Mediator complex. This gene is located within the Smith-Magenis syndrome region on chromosome 17. [provided by RefSeq]

Other Designations

mediator of RNA polymerase II transcription, subunit 9 homolog|mediator subunit 25