


MED29 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # H00055588-T03


Size 100 uL

Applications



SDS-PAGE Gel

MED29 transfected lysate.



Western Blot

Lane 1: MED29 transfected lysate (23.50 KDa)

Lane 2: Non-transfected lysate.

Specification

Transfected Cell Line	293T
Plasmid	pCMV-MED29 full-length
Host	Human
Theoretical MW (kDa)	23.5
Interspecies Antigen Sequence	Mouse (95); Rat (96)

Quality Control Testing

Transient overexpression cell lysate was tested with Anti-MED29 antibody ([H00055588-B01P](#)) by Western Blots.
SDS-PAGE Gel
MED29 transfected lysate.
Western Blot
Lane 1: MED29 transfected lysate (23.50 KDa)
Lane 2: Non-transfected lysate.

Storage Buffer

1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction

Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot

Gene Info — MED29

Entrez GeneID

[55588](#)

GeneBank Accession#

[NM_017592](#)

Protein Accession#

[NP_060062](#)

Gene Name

MED29

Gene Alias

DKFZp434H247, IXL

Gene Description

mediator complex subunit 29

Gene Ontology

[Hyperlink](#)

Gene Summary

MED29 is a subunit of the Mediator complex, a multiprotein coactivator of RNA transcription that interacts with DNA-bound transcriptional activators, RNA polymerase II (see MIM 180660), and general initiation factors (Sato et al., 2003 [PubMed 14576168]).[supplied by OMIM]

Other Designations

intersex-like