

# PCDHGA10 rabbit monoclonal antibody

Catalog # H00056106-K      Size 100 ug x up to 3

## Specification

<b>Product Description</b>	Rabbit monoclonal antibody raised against a human PCDHGA10 peptide using ARM Technology.
<b>Immunogen</b>	A synthetic peptide of human PCDHGA10 is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
<b>Host</b>	Rabbit
<b>Library Construction</b>	Non-fusion antibody library from rabbit spleen ( <a href="#">ARM Technology</a> ).
<b>Expression</b>	Overexpression vector and transfection into 293H cell line.
<b>Reactivity</b>	Human
<b>Purification</b>	Protein A
<b>Isotype</b>	IgG
<b>Quality Control Testing</b>	Antibody reactive against human PCDHGA10 peptide by ELISA and mammalian transfected lysate by Western Blot.
<b>Storage Buffer</b>	In 1x PBS, pH 7.4
<b>Storage Instruction</b>	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
<b>Deliverable</b>	Up to three rabbit IgG clones of 100 ug each will be delivered to customer.
<b>Note</b>	1. Customer may provide cell or tissue lysate for antibody screening. 2. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering including F(ab) <sub>2</sub> , IgG, scFv and different Fc and non-Fc conjugates per customer request.

## Applications

- Western Blot (Transfected lysate)

[Protocol Download](#)

- ELISA

## Gene Info — PCDHGA10

Entrez GeneID [56106](#)

GeneBank Accession# [PCDHGA10](#)

Gene Name PCDHGA10

Gene Alias PCDH-GAMMA-A10

Gene Description protocadherin gamma subfamily A, 10

Omim ID [606297](#)

Gene Ontology [Hyperlink](#)

**Gene Summary**

This gene is a member of the protocadherin gamma gene cluster, one of three related clusters tandemly linked on chromosome five. These gene clusters have an immunoglobulin-like organization, suggesting that a novel mechanism may be involved in their regulation and expression. The gamma gene cluster includes 22 genes divided into 3 subfamilies. Subfamily A contains 12 genes, subfamily B contains 7 genes and 2 pseudogenes, and the more distantly related subfamily C contains 3 genes. The tandem array of 22 large, variable region exons are followed by a constant region, containing 3 exons shared by all genes in the cluster. Each variable region exon encodes the extracellular region, which includes 6 cadherin ectodomains and a transmembrane region. The constant region exons encode the common cytoplasmic region. These neural cadherin-like cell adhesion proteins most likely play a critical role in the establishment and function of specific cell-cell connections in the brain. Alternative splicing has been described for the gamma cluster genes. [provided by RefSeq]

Other Designations -