


PCDHGA10 (Human) Recombinant Protein (Q01)

Catalog # H00056106-Q01

Size 10 ug, 25 ug

Applications



Specification

Product Description	Human PCDHGA10 partial ORF (NP_061736.1, 219 a.a. - 316 a.a.) recombinant protein with GST-tag at N-terminal.
Sequence	SDGGDPLRSGTVLVSVTVFDANDNAPVFTLPEYRVSVPENLPVGTQLLTVTATDRDEGANGEVT YSFRKLDPDTQLLKFKQLNKYTGEIKISENLDYEET
Host	Wheat Germ (in vitro)
Theoretical MW (kDa)	36.52
Interspecies Antigen Sequence	Mouse (85); Rat (84)
Preparation Method	in vitro wheat germ expression system
Purification	Glutathione Sepharose 4 Fast Flow
Quality Control Testing	12.5% SDS-PAGE Stained with Coomassie Blue.
Storage Buffer	50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.
Storage Instruction	Store at -80°C. Aliquot to avoid repeated freezing and thawing.
Note	Best use within three months from the date of receipt of this protein.

Applications

- Enzyme-linked Immunoabsorbent Assay
- Western Blot (Recombinant protein)
- Antibody Production
- Protein Array

Gene Info — PCDHGA10

Entrez GeneID [56106](#)

GeneBank Accession# [NM_018913](#)

Protein Accession# [NP_061736.1](#)

Gene Name PCDHGA10

Gene Alias PCDH-GAMMA-A10

Gene Description protocadherin gamma subfamily A, 10

Omim ID [606297](#)

Gene Ontology [Hyperlink](#)

Gene Summary

This gene is a member of the protocadherin gamma gene cluster, one of three related clusters tandemly linked on chromosome five. These gene clusters have an immunoglobulin-like organization, suggesting that a novel mechanism may be involved in their regulation and expression. The gamma gene cluster includes 22 genes divided into 3 subfamilies. Subfamily A contains 12 genes, subfamily B contains 7 genes and 2 pseudogenes, and the more distantly related subfamily C contains 3 genes. The tandem array of 22 large, variable region exons are followed by a constant region, containing 3 exons shared by all genes in the cluster. Each variable region exon encodes the extracellular region, which includes 6 cadherin ectodomains and a transmembrane region. The constant region exons encode the common cytoplasmic region. These neural cadherin-like cell adhesion proteins most likely play a critical role in the establishment and function of specific cell-cell connections in the brain. Alternative splicing has been described for the gamma cluster genes. [provided by RefSeq]

Other Designations -