
## PCDHGA3 (Human) Recombinant Protein (Q01)

Catalog # H00056112-Q01 Size 10 ug, 25 ug

## Applications



Specification	
Product Description	Human PCDHGA3 partial ORF ( NP_061739, 98 a.a 206 a.a.) recombinant protein with GST-tag at N-terminal.
Sequence	QIPLCLVKINILVEDKLKIFEVEIEIKDINDNAPNFPTEELEIKIGELTVPGTRFPIKTAFDPDVGINSLQN YKLSPNDYFSLAVNSVSEGAKYPELVLERALDREKKE
Host	Wheat Germ (in vitro)
Theoretical MW (kDa)	37.73
Interspecies Antigen Sequence	Mouse (85); Rat (86)
Preparation Method	in vitro wheat germ expression system
Purification	Glutathione Sepharose 4 Fast Flow
Quality Control Testing	12.5% SDS-PAGE Stained with Coomassie Blue.
Storage Buffer	50 mM Tris-HCI, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.
Storage Instruction	Store at -80°C. Aliquot to avoid repeated freezing and thawing.
Note	Best use within three months from the date of receipt of this protein.



## Applications

- Enzyme-linked Immunoabsorbent Assay
- Western Blot (Recombinant protein)
- Antibody Production
- Protein Array

Gene Info — PCDHGA3	
Entrez GenelD	<u>56112</u>
GeneBank Accession#	<u>NM_018916</u>
Protein Accession#	<u>NP_061739</u>
Gene Name	PCDHGA3
Gene Alias	PCDH-GAMMA-A3
Gene Description	protocadherin gamma subfamily A, 3
Omim ID	<u>606290</u>
Gene Ontology	<u>Hyperlink</u>
Gene Summary	This gene is a member of the protocadherin gamma gene cluster, one of three related clusters tan demly linked on chromosome five. These gene clusters have an immunoglobulin-like organization, suggesting that a novel mechanism may be involved in their regulation and expression. The gam ma gene cluster includes 22 genes divided into 3 subfamilies. Subfamily A contains 12 genes, su bfamily B contains 7 genes and 2 pseudogenes, and the more distantly related subfamily C contains 3 genes. The tandem array of 22 large, variable region exons are followed by a constant region, containing 3 exons shared by all genes in the cluster. Each variable region exon encodes the ext racellular region, which includes 6 cadherin ectodomains and a transmembrane region. The const
	ant region exons encode the common cytoplasmic region. These neural cadherin-like cell adhesio n proteins most likely play a critical role in the establishment and function of specific cell-cell conn ections in the brain. Alternative splicing has been described for the gamma cluster genes. [provid ed by RefSeq



**Product Information** 

• Tobacco Use Disorder