


## PCDHB15 monoclonal antibody (M02), clone 2E10

Catalog # H00056121-M02 Size 100 ug

## Applications



Western Blot detection against Immunogen (36.74 KDa) .

Specification	
Product Description	Mouse monoclonal antibody raised against a partial recombinant PCDHB15.
Immunogen	PCDHB15 (NP_061758, 207 a.a. ~ 306 a.a) partial recombinant protein with GST tag. MW of the G ST tag alone is 26 KDa.
Sequence	ELRLTLTAVDGGSPPRSGTVQILILVLDANDNAPEFVQALYEVQVPENSPVGSLVVKVSARDLDT GTNGEISYSLYYSSQEIDKPFELSSLSGEIRLIKK
Host	Mouse
Reactivity	Human
Interspecies Antigen Sequence	Mouse (76); Rat (76)
Isotype	lgG2a Kappa
Quality Control Testing	Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.74 KDa) .
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.



## Applications

- Western Blot (Recombinant protein)
  <u>Protocol Download</u>
- ELISA

## Gene Info — PCDHB15

Entrez GenelD	<u>56121</u>
GeneBank Accession#	<u>NM_018935</u>
Protein Accession#	<u>NP_061758</u>
Gene Name	PCDHB15
Gene Alias	PCDH-BETA15
Gene Description	protocadherin beta 15
Omim ID	<u>606341</u>
Gene Ontology	Hyperlink
Gene Summary	This gene is a member of the protocadherin beta gene cluster, one of three related gene clusters t andemly linked on chromosome five. The gene clusters demonstrate an unusual genomic organiz ation similar to that of B-cell and T-cell receptor gene clusters. The beta cluster contains 16 genes and 3 pseudogenes, each encoding 6 extracellular cadherin domains and a cytoplasmic tail that d eviates from others in the cadherin superfamily. The extracellular domains interact in a homophilic manner to specify differential cell-cell connections. Unlike the alpha and gamma clusters, the trans cripts from these genes are made up of only one large exon, not sharing common 3' exons as exp ected. These neural cadherin-like cell adhesion proteins are integral plasma membrane proteins. Their specific functions are unknown but they most likely play a critical role in the establishment an d function of specific cell-cell neural connections. [provided by RefSeq
Other Designations	-