PCDHB13 rabbit monoclonal antibody

Size

Catalog # H00056123-K

100 ug x up to 3

Specification **Product Description** Rabbit monoclonal antibody raised against a human PCDHB13 peptide using ARM Technology. Immunogen A synthetic peptide of human PCDHB13 is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence. Host Rabbit Library Construction Non-fusion antibody library from rabbit spleen (ARM Technology). Expression Overexpression vector and transfection into 293H cell line. Reactivity Human **Purification** Protein A lsotype lgG **Quality Control Testing** Antibody reactive against human PCDHB13 peptide by ELISA and mammalian transfected lysate by Western Blot. **Storage Buffer** In 1x PBS, pH 7.4 **Storage Instruction** Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing. Deliverable Up to three rabbit IgG clones of 100 ug each will be delivered to customer. Note 1. Customer may provide cell or tissue lysate for antibody screening. 2. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering in cluding F(ab)₂, IgG, scFv and different Fc and non-Fc conjugates per customer request.

Applications

• Western Blot (Transfected lysate)

Protocol Download

• ELISA

Gene Info — PCDHB13	
Entrez GenelD	<u>56123</u>
GeneBank Accession#	PCDHB13
Gene Name	PCDHB13
Gene Alias	PCDH-BETA13
Gene Description	protocadherin beta 13
Omim ID	<u>606339</u>
Gene Ontology	Hyperlink
Gene Summary	This gene is a member of the protocadherin beta gene cluster, one of three related gene clusters t andemly linked on chromosome five. The gene clusters demonstrate an unusual genomic organiz ation similar to that of B-cell and T-cell receptor gene clusters. The beta cluster contains 16 genes and 3 pseudogenes, each encoding 6 extracellular cadherin domains and a cytoplasmic tail that d eviates from others in the cadherin superfamily. The extracellular domains interact in a homophilic manner to specify differential cell-cell connections. Unlike the alpha and gamma clusters, the trans cripts from these genes are made up of only one large exon, not sharing common 3' exons as exp ected. These neural cadherin-like cell adhesion proteins are integral plasma membrane proteins. Their specific functions are unknown but they most likely play a critical role in the establishment an d function of specific cell-cell neural connections. [provided by RefSeq
Other Designations	-