


Full-Length

PCDHB13 (Human) Recombinant Protein (P01)

Catalog # H00056123-P01

Size 25 ug, 10 ug

Applications



Specification

Product Description

Human PCDHB13 full-length ORF (NP_061756.1, 1 a.a. - 798 a.a.) recombinant protein with GST-tag at N-terminal.

Sequence

MEASGKLCRQRQVLFSFLLLGLSLAGAAEPRSYSVVEETEGSSFVTNLAKDLGLEQREFSRRGV
RVVSRGNKLHLQLNQETADLLLNEKLDREDLCGHTEPCVLRQVLLESPFEFFQAEQVIDINDHS
PVFLDKQMLVKVSESSPPGTTFFPLKNAEDLDVGQNNIENYIISPNSYFRVLTRKRS DGRKYPELVL
DKALDREEEAELRLTLTALDGGSPPRSGTAQVYIEVLDVNDNAPEFEQPFYRVQISEDSPVGFLV
VKVSATDVDVTGVNGEISYSLFQASEEIGKTFKINPLTGEIELKKQLDFEKLQSYEVNIEARDAGTFS
GKCTVLIQVIDVNDHAPEVTMSAFTSPIENAPETVVALFSVSDLDSENGKISCSIQEDLPFLLKS
AENFYTLTERPLDRESRAEYNITITVDLGTPLMLITQLNMTVLIADVNDNAPAFQTQSYTLFVRENNS
PALHIRSVSATDRDSGTNAQVTYSLPPQDPLPLTSLVSINADNGHLFALRSLDYEALQGFQFRV
GASDHGSPALSSEALVRVVLDANDNSPFVLYPLQNGSAPCTELVPRAAEPGYLVTKVAVDVG
DSGQNAWLSYQLLKATELGFLGVWAHNGEVRTARLLSERDAAKHRLVVLVKDNGEPPRSATATL
HVLLVDGFSQPYLPLPEAAPTQAQADLLTVYLVALASVSSLFLFSVLLFVAVRLCRRSRAASVG
RCLVPEGPLPGHLVDMMSGTRTSLSQSYQYEVCLAGGSGTNEFKFLKPIIPNFPFPQCPGKEIQGNSTF
PNNFGFNIQ

Host

Wheat Germ (in vitro)

Theoretical MW (kDa)

114

Preparation Method

[in vitro wheat germ expression system](#)

Purification

Glutathione Sepharose 4 Fast Flow

Quality Control Testing	12.5% SDS-PAGE Stained with Coomassie Blue.
Storage Buffer	50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.
Storage Instruction	Store at -80°C. Aliquot to avoid repeated freezing and thawing.
Note	Best use within three months from the date of receipt of this protein.

Applications

- Enzyme-linked Immunoabsorbent Assay
- Western Blot (Recombinant protein)
- Antibody Production
- Protein Array

Gene Info — PCDHB13

Entrez GeneID	56123
GeneBank Accession#	NM_018933.2
Protein Accession#	NP_061756.1
Gene Name	PCDHB13
Gene Alias	PCDH-BETA13
Gene Description	protocadherin beta 13
Omim ID	606339
Gene Ontology	Hyperlink

Gene Summary

This gene is a member of the protocadherin beta gene cluster, one of three related gene clusters tandemly linked on chromosome five. The gene clusters demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The beta cluster contains 16 genes and 3 pseudogenes, each encoding 6 extracellular cadherin domains and a cytoplasmic tail that deviates from others in the cadherin superfamily. The extracellular domains interact in a homophilic manner to specify differential cell-cell connections. Unlike the alpha and gamma clusters, the transcripts from these genes are made up of only one large exon, not sharing common 3' exons as expected. These neural cadherin-like cell adhesion proteins are integral plasma membrane proteins. Their specific functions are unknown but they most likely play a critical role in the establishment and function of specific cell-cell neural connections. [provided by RefSeq]

Other Designations

-