

## PCDHB7 rabbit monoclonal antibody

Catalog # H00056129-K      Size 100 ug x up to 3

### Specification

<b>Product Description</b>	Rabbit monoclonal antibody raised against a human PCDHB7 peptide using ARM Technology.
<b>Immunogen</b>	A synthetic peptide of human PCDHB7 is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
<b>Host</b>	Rabbit
<b>Library Construction</b>	Non-fusion antibody library from rabbit spleen ( <a href="#">ARM Technology</a> ).
<b>Expression</b>	Overexpression vector and transfection into 293H cell line.
<b>Reactivity</b>	Human
<b>Purification</b>	Protein A
<b>Isotype</b>	IgG
<b>Quality Control Testing</b>	Antibody reactive against human PCDHB7 peptide by ELISA and mammalian transfected lysate by Western Blot.
<b>Storage Buffer</b>	In 1x PBS, pH 7.4
<b>Storage Instruction</b>	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
<b>Deliverable</b>	Up to three rabbit IgG clones of 100 ug each will be delivered to customer.
<b>Note</b>	1. Customer may provide cell or tissue lysate for antibody screening. 2. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering including F(ab) <sub>2</sub> , IgG, scFv and different Fc and non-Fc conjugates per customer request.

### Applications

- Western Blot (Transfected lysate)

[Protocol Download](#)

- ELISA

## Gene Info — PCDHB7

Entrez GeneID	<a href="#">56129</a>
GeneBank Accession#	<a href="#">PCDHB7</a>
Gene Name	PCDHB7
Gene Alias	MGC111391, MGC163205, PCDH-BETA7
Gene Description	protocadherin beta 7
Omim ID	<a href="#">606333</a>
Gene Ontology	<a href="#">Hyperlink</a>
Gene Summary	<p>This gene is a member of the protocadherin beta gene cluster, one of three related gene clusters tandemly linked on chromosome five. The gene clusters demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The beta cluster contains 16 genes and 3 pseudogenes, each encoding 6 extracellular cadherin domains and a cytoplasmic tail that deviates from others in the cadherin superfamily. The extracellular domains interact in a homophilic manner to specify differential cell-cell connections. Unlike the alpha and gamma clusters, the transcripts from these genes are made up of only one large exon, not sharing common 3' exons as expected. These neural cadherin-like cell adhesion proteins are integral plasma membrane proteins. Their specific functions are unknown but they most likely play a critical role in the establishment and function of specific cell-cell neural connections. The transcript for this particular family member uses more than one polyadenylation site. [provided by RefSeq]</p>
Other Designations	-