


Full-Length

PCDHB2 (Human) Recombinant Protein (P01)

Catalog # H00056133-P01

Size 25 ug, 10 ug

Applications



Specification

Product Description

Human PCDHB2 full-length ORF (AAH98575.1, 1 a.a. - 798 a.a.) recombinant protein with GST-tag at N-terminal.

Sequence

MEAGEGKERVPKQRQVLFFVLLGIAQASCQPRHYSVAEETESGSFVANLLKDLGLEIGELAVRG
ARVVSCKGKKMHLQFDRQTGDLLEKLDREELCGPTEPCVLPFQVLLENPLQFFQAE LRIRDVN
DHSPVFLDKEILLKIPESITPGTTFLIERAQDLVDGTNSLQNYTISPNFHFHLNLQDSL DGIILPQLVLN
RALDREEQPEIRLTALTALDGGSPPRSGTALVRIEVDINDNVPEFAKLLYEVQIPEDSPVGSQVAM
SARDLDIGTNGEISYAFSQASEDIRKTFRLSAKSGELLLRQKLDFESIQT YTVNIQATDGGGLSGTCV
VFVQVMDLNDNPPELTMSTLINQIPENLQDTLAVFSVSDPDSDGNRMVCSIQDDL PFFLKPSVE
NFYTLVISTALDRETRSEYNITITVDFGTPRLKTEHNITVLVSDVNDNAPFTQTSYTLFVRENNSPA
LHIGSVSATDRDSGTNAQVTYSLPPQDPHPLASLV SINADNGHLFALQSLDYEALQAFEFVRGA
ADRGSPALSSEALVRVLVDANDNSPFVLYPLQNGSAPCTELVPRAAEPGYLVTKVVAVDGDSG
QNAWLSYQLLKATEPGLFGVWAHNGEVRTARLLRERDAAKQRLVVLVKDNGEPPRSATATLHVL
LVDGFSQPYLLPEAAPAQADLLTVYLVVALASVSSLFLFSVLLFVAVRLCRRSRAASVGRCS
VPEGPFPQMVDVSGTGTLSQSYQYEVCLTGESGTNEFKFLKPIIPNFVAQGAERVSEANPSFRK
SFEFT

Host

Wheat Germ (in vitro)

Theoretical MW (kDa)

113.7

Interspecies Antigen Sequence

Mouse (76)

Preparation Method	in vitro wheat germ expression system
Purification	Glutathione Sepharose 4 Fast Flow
Quality Control Testing	12.5% SDS-PAGE Stained with Coomassie Blue.
Storage Buffer	50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.
Storage Instruction	Store at -80°C. Aliquot to avoid repeated freezing and thawing.
Note	Best use within three months from the date of receipt of this protein.

Applications

- Enzyme-linked Immunoabsorbent Assay
- Western Blot (Recombinant protein)
- Antibody Production
- Protein Array

Gene Info — PCDHB2

Entrez GeneID	56133
GeneBank Accession#	BC098575.1
Protein Accession#	AAH98575.1
Gene Name	PCDHB2
Gene Alias	MGC111392, PCDH-BETA2
Gene Description	protocadherin beta 2
Omim ID	606328
Gene Ontology	Hyperlink

Gene Summary

This gene is a member of the protocadherin beta gene cluster, one of three related gene clusters tandemly linked on chromosome five. The gene clusters demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The beta cluster contains 16 genes and 3 pseudogenes, each encoding 6 extracellular cadherin domains and a cytoplasmic tail that deviates from others in the cadherin superfamily. The extracellular domains interact in a homophilic manner to specify differential cell-cell connections. Unlike the alpha and gamma clusters, the transcripts from these genes are made up of only one large exon, not sharing common 3' exons as expected. These neural cadherin-like cell adhesion proteins are integral plasma membrane proteins. Their specific functions are unknown but they most likely play a critical role in the establishment and function of specific cell-cell neural connections. [provided by RefSeq]

Other Designations

-