WDR18 rabbit monoclonal antibody

Catalog # H00057418-K

Size 100 ug x up to 3

Specification **Product Description** Rabbit monoclonal antibody raised against a human WDR18 peptide using ARM Technology. Immunogen A synthetic peptide of human WDR18 is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence. Host Rabbit Library Construction Non-fusion antibody library from rabbit spleen (ARM Technology). Expression Overexpression vector and transfection into 293H cell line. Reactivity Human **Purification** Protein A lsotype lgG **Quality Control Testing** Antibody reactive against human WDR18 peptide by ELISA and mammalian transfected lysate by W estern Blot. **Storage Buffer** In 1x PBS, pH 7.4 **Storage Instruction** Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing. Deliverable Up to three rabbit IgG clones of 100 ug each will be delivered to customer. Note 1. Customer may provide cell or tissue lysate for antibody screening. 2. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering in cluding F(ab)₂, IgG, scFv and different Fc and non-Fc conjugates per customer request.

Applications

Western Blot (Transfected lysate)

Protocol Download

• ELISA

Gene Info — WDR18

ed regions of approximately 40 amino acids typically bracketed by gly-his and trp-asp (GH-WD), which may facilitate formation of heterotrimeric or multiprotein complexes. Members of this family		
Gene Name WDR18 Gene Alias MGC2436, R32184_1 Gene Description WD repeat domain 18 Gene Ontology Hyperlink Gene Summary This gene encodes a member of the WD repeat protein family. WD repeats are minimally conserted regions of approximately 40 amino acids typically bracketed by gly-his and trp-asp (GH-WD), which may facilitate formation of heterotrimeric or multiprotein complexes. Members of this family are involved in a variety of cellular processes, including cell cycle progression, signal transduction apoptosis, and gene regulation. [provided by RefSeq	Entrez GenelD	57418
Gene Alias MGC2436, R32184_1 Gene Description WD repeat domain 18 Gene Ontology Hyperlink Gene Summary This gene encodes a member of the WD repeat protein family. WD repeats are minimally consered regions of approximately 40 amino acids typically bracketed by gly-his and trp-asp (GH-WD), which may facilitate formation of heterotrimeric or multiprotein complexes. Members of this family are involved in a variety of cellular processes, including cell cycle progression, signal transduction apoptosis, and gene regulation. [provided by RefSeq	GeneBank Accession#	<u>WDR18</u>
Gene Description WD repeat domain 18 Gene Ontology Hyperlink Gene Summary This gene encodes a member of the WD repeat protein family. WD repeats are minimally conser ed regions of approximately 40 amino acids typically bracketed by gly-his and trp-asp (GH-WD), which may facilitate formation of heterotrimeric or multiprotein complexes. Members of this family are involved in a variety of cellular processes, including cell cycle progression, signal transduction apoptosis, and gene regulation. [provided by RefSeq	Gene Name	WDR18
Gene Ontology Hyperlink Gene Summary This gene encodes a member of the WD repeat protein family. WD repeats are minimally consered regions of approximately 40 amino acids typically bracketed by gly-his and trp-asp (GH-WD), which may facilitate formation of heterotrimeric or multiprotein complexes. Members of this family are involved in a variety of cellular processes, including cell cycle progression, signal transduction apoptosis, and gene regulation. [provided by RefSeq	Gene Alias	MGC2436, R32184_1
Gene Summary This gene encodes a member of the WD repeat protein family. WD repeats are minimally conser ed regions of approximately 40 amino acids typically bracketed by gly-his and trp-asp (GH-WD), which may facilitate formation of heterotrimeric or multiprotein complexes. Members of this family are involved in a variety of cellular processes, including cell cycle progression, signal transduction apoptosis, and gene regulation. [provided by RefSeq	Gene Description	WD repeat domain 18
ed regions of approximately 40 amino acids typically bracketed by gly-his and trp-asp (GH-WD), which may facilitate formation of heterotrimeric or multiprotein complexes. Members of this family are involved in a variety of cellular processes, including cell cycle progression, signal transduction apoptosis, and gene regulation. [provided by RefSeq	Gene Ontology	<u>Hyperlink</u>
Other Designations -	Gene Summary	which may facilitate formation of heterotrimeric or multiprotein complexes. Members of this family are involved in a variety of cellular processes, including cell cycle progression, signal transduction,
	Other Designations	-