


CIDEC 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # H00063924-T01


Size 100 uL

Applications



SDS-PAGE Gel

CIDEc transfected lysate



Western Blot

Lane 1: CIDEc transfected lysate (26.29 KDa).

Lane 2: Non-transfected lysate.

Specification

Product Description

Transfected Cell Line	293T
Plasmid	pCMV-CIDE-3 full-length
Host	Human
Theoretical MW (kDa)	26.29
Interspecies Antigen Sequence	Mouse (79); Rat (79)

Quality Control Testing

Transient overexpression cell lysate was tested with Anti-CIDE-3 antibody ([H00063924-B01](#)) by Western Blots.
SDS-PAGE Gel
CIDE-3 transfected lysate
Western Blot
Lane 1: CIDE-3 transfected lysate (26.29 KDa).
Lane 2: Non-transfected lysate.

Storage Buffer

1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction

Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot

Gene Info — CIDE-3

Entrez GeneID[63924](#)**GeneBank Accession#**[BC016851](#)**Protein Accession#**[AAH16851](#)**Gene Name**

CIDE-3

Gene Alias

CIDE-3, FLJ20871, Fsp27

Gene Description

cell death-inducing DFFA-like effector c

Gene Ontology[Hyperlink](#)**Gene Summary**

DNA fragmentation factor (DFF) induces the fragmentation of DNA associated with apoptosis. A novel family of cell death-inducing DFF45 (MIM 601882)-like effectors (CIDEs), including CIDE-3, can also promote apoptosis (Liang et al., 2003 [PubMed 12429024]).[supplied by OMIM]

Other Designations

cell death activator CIDE-3|fat specific protein 27