


RNF25 293T Cell Transient Overexpression Lysate(Denatured)


Catalog # H00064320-T01 Size 100 uL

Applications



SDS-PAGE Gel

RNF25 transfected lysate.



Western Blot

Lane 1: RNF25 transfected lysate (50.6 KDa)

Lane 2: Non-transfected lysate.

| Specification | |
|----------------------------------|------------------------|
| Transfected Cell Line | 293T |
| Plasmid | pCMV-RNF25 full-length |
| Host | Human |
| Theoretical MW (kDa) | 50.6 |
| Interspecies Antigen Sequence | Mouse (82); Rat (84) |



Product Information

| Quality Control Testing | Transient overexpression cell lysate was tested with Anti-RNF25 antibody (H00064320-B01) by Wes tern Blots. SDS-PAGE Gel RNF25 transfected lysate. Western Blot Lane 1: RNF25 transfected lysate (50.6 KDa) Lane 2: Non-transfected lysate. |
|-------------------------|---|
| Storage Buffer | 1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bro mophenol blue) |
| Storage Instruction | Store at -80°C. Aliquot to avoid repeated freezing and thawing. |

Applications

Western Blot

| Gene Info — RNF25 | |
|---------------------|---|
| Entrez GenelD | <u>64320</u> |
| GeneBank Accession# | NM_022453.2 |
| Protein Accession# | NP_071898.2 |
| Gene Name | RNF25 |
| Gene Alias | AO7, FLJ13906 |
| Gene Description | ring finger protein 25 |
| Gene Ontology | <u>Hyperlink</u> |
| Gene Summary | The protein encoded by this gene contains a RING finger motif. The mouse counterpart of this prot ein has been shown to interact with Rela, the p65 subunit of NF-kappaB (NFKB), and modulate N FKB-mediated transcription activity. The mouse protein also binds ubiquitin-conjugating enzymes (E2s) and is a substrate for E2-dependent ubiquitination. [provided by RefSeq |
| Other Designations | ring finger protein AO7 |