


MaxPab®

FSD1 purified MaxPab rabbit polyclonal antibody (D01P)

Catalog # H00079187-D01P

Size 100 ug

Applications



Western Blot (Transfected lysate)

Western Blot analysis of FSD1 expression in transfected 293T cell line ([H00079187-T01](#)) by FSD1 MaxPab polyclonal antibody.

Lane 1: FSD1 transfected lysate(55.80 kDa).

Lane 2: Non-transfected lysate.

Specification

Product Description	Rabbit polyclonal antibody raised against a full-length human FSD1 protein.
Immunogen	FSD1 (NP_077309.1, 1 a.a. ~ 496 a.a) full-length human protein.
Sequence	MEEQREALRKIIKTLAVKNEEIQSFIYSLKQMLLNVEANSAKVQEDLEAEFQSLFSLLEELKEGMLM KIKQDRASRTYELQNQLAACTRALESSEELLETANQTLQAMDSEDFPQAAKQIKDGVTMAPAFRL SLKAKVSDNMSHLMVDFAQERQMLQALKFLPVPSAPVIDLAESLVADNCVTLVWRMPDEDSKID HYVLEYRRTNFEGPPRLKEDQPWMVIEGIRQTEYTLTGLKFDMKYMNFRVKACNKAVAGEFSEPV TLETPAFMFRLDASTSHQNLRVDDLSEWDAMGGKVQDIKAREKDGGKRTASPINSPARGTPSP KRMPSGRGRDRFTAESYTVLGDTLIDGGEHYWEVRYEPDSKAFGVGVAYRSLGRFEQLGKTAA SWCLHVNWLQVSFTAKHANKVKVLDAPVPDCLGVHCDHFHQGLLSFYNARTKQVLHTFKTRFT QPLLPAFTVWCGSFQVTTGLQVPSAVRCLQKRGSATSSSNTSLT
Host	Rabbit
Reactivity	Human
Interspecies Antigen Sequence	Mouse (93)
Quality Control Testing	Antibody reactive against mammalian transfected lysate.

Storage Buffer

In 1x PBS, pH 7.4

Storage Instruction

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Transfected lysate)

Western Blot analysis of FSD1 expression in transfected 293T cell line ([H00079187-T01](#)) by FSD1 MaxPab polyclonal antibody.

Lane 1: FSD1 transfected lysate(55.80 KDa).

Lane 2: Non-transfected lysate.

[Protocol Download](#)

Gene Info — FSD1

Entrez GeneID[79187](#)**GeneBank Accession#**[NM_024333.1](#)**Protein Accession#**[NP_077309.1](#)**Gene Name**

FSD1

Gene Alias

GLFND, MGC3213, MIR1

Gene Description

fibronectin type III and SPRY domain containing 1

Omim ID[609828](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

This gene encodes a centrosome associated protein that is characterized by an N-terminal coiled-coil region downstream of B-box (BBC) domain, a central fibronectin type III domain, and a C-terminal repeats in splA and RyR (SPRY) domain. The encoded protein associates with a subset of microtubules and may be involved in the stability and organization of microtubules during cytokinesis. [provided by RefSeq]

Other Designations

fibronectin type 3 and SPRY (splA, ryanodine) domain containing (with coiled-coil motif) 1|fibronectin type 3 and SPRY domain containing 1|fibronectin type 3 and SPRY domain-containing protein