


HDAC11 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # H00079885-T01


Size 100 uL

Applications



SDS-PAGE Gel

HDAC11 transfected lysate.



Western Blot

Lane 1: HDAC11 transfected lysate (38.28 KDa)

Lane 2: Non-transfected lysate.

Specification

Transfected Cell Line	293T
Plasmid	pCMV-HDAC11 full-length
Host	Human
Theoretical MW (kDa)	38.28
Interspecies Antigen Sequence	Mouse (92); Rat (91)

Quality Control Testing

Transient overexpression cell lysate was tested with Anti-HDAC11 antibody ([H00079885-B01](#)) by Western Blots.
SDS-PAGE Gel
HDAC11 transfected lysate.
Western Blot
Lane 1: HDAC11 transfected lysate (38.28 KDa)
Lane 2: Non-transfected lysate.

Storage Buffer

1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction

Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot

Gene Info — HDAC11

Entrez GeneID[79885](#)**GeneBank Accession#**[BC009676.1](#)**Protein Accession#**[AAH09676.1](#)**Gene Name**

HDAC11

Gene Alias

FLJ22237

Gene Description

histone deacetylase 11

Omim ID[607226](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

This gene encodes a class IV histone deacetylase. The encoded protein is localized to the nucleus and may be involved in regulating the expression of interleukin 10. Alternative splicing results in multiple transcript variants

Other Designations

-