


NETO2 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # H00081831-T01


Size 100 uL

Applications



SDS-PAGE Gel

NETO2 transfected lysate.



Western Blot

Lane 1: NETO2 transfected lysate (17.00 KDa)

Lane 2: Non-transfected lysate.

Specification

Transfected Cell Line 293T

Plasmid pCMV-NETO2 full-length

Host Human

Theoretical MW (kDa) 17

Quality Control Testing Transient overexpression cell lysate was tested with Anti-NETO2 antibody ([H00081831-B01P](#)) by Western Blots.
 SDS-PAGE Gel
 NETO2 transfected lysate.
 Western Blot
 Lane 1: NETO2 transfected lysate (17.00 KDa)
 Lane 2: Non-transfected lysate.

Storage Buffer

1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction

Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot

Gene Info — NETO2

Entrez GeneID[81831](#)**GeneBank Accession#**[BC012381.1](#)**Protein Accession#**[AAH12381.1](#)**Gene Name**

NETO2

Gene Alias

FLJ10430, FLJ14724, FLJ90456, NEOT2

Gene Description

neuropilin (NRP) and tolloid (TLL)-like 2

Omim ID[607974](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

This gene encodes a predicted transmembrane protein containing two extracellular CUB domains followed by a low-density lipoprotein class A (LDLa) domain. It also has an intracellular FXNPXY-like motif, which has been shown in other proteins to be essential for the internalization of clathrin-coated pits during endocytosis. Alternatively spliced transcript variants have been observed, but they have not been fully characterized. [provided by RefSeq]

Other Designations

neuropilin- and tolloid-like protein 2