


MKI67IP 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # H00084365-T01


Size 100 uL

Applications



SDS-PAGE Gel

MKI67IP transfected lysate.



Western Blot

Lane 1: MKI67IP transfected lysate (34.3 KDa)

Lane 2: Non-transfected lysate.

Specification

Transfected Cell Line	293T
Plasmid	pCMV-MKI67IP full-length
Host	Human
Theoretical MW (kDa)	34.3
Interspecies Antigen Sequence	Mouse (56); Rat (59)

Quality Control Testing

Transient overexpression cell lysate was tested with Anti-MKI67IP antibody ([H00084365-B01](#)) by Western Blots.
SDS-PAGE Gel
MKI67IP transfected lysate.
Western Blot
Lane 1: MKI67IP transfected lysate (34.3 KDa)
Lane 2: Non-transfected lysate.

Storage Buffer

1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction

Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot

Gene Info — MKI67IP

Entrez GeneID

[84365](#)

GeneBank Accession#

[NM_032390.3](#)

Protein Accession#

-

Gene Name

MKI67IP

Gene Alias

NIFK, Nopp34

Gene Description

MKI67 (FHA domain) interacting nucleolar phosphoprotein

Gene Ontology

[Hyperlink](#)

Gene Summary

This gene encodes a protein that interacts with the forkhead-associated domain of the Ki-67 antigen. The encoded protein may bind RNA and may play a role in mitosis and cell cycle progression. Multiple pseudogenes exist on chromosomes 5, 10, 12, 15, and 19.[provided by RefSeq]

Other Designations

nucleolar phosphoprotein Nopp34|nucleolar protein interacting with the FHA domain of pKi-67