


IL17RC 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # H00084818-T01


Size 100 uL

Applications



SDS-PAGE Gel

IL17RC transfected lysate.



Western Blot

Lane 1: IL17RC transfected lysate (59.29 KDa)

Lane 2: Non-transfected lysate.

Specification

Transfected Cell Line 293T

Plasmid pCMV-IL17RC full-length

Host Human

Theoretical MW (kDa) 59.29

Quality Control Testing Transient overexpression cell lysate was tested with Anti-IL17RC antibody ([H00084818-B01](#)) by Western Blots.
 SDS-PAGE Gel
 IL17RC transfected lysate.
 Western Blot
 Lane 1: IL17RC transfected lysate (59.29 KDa)
 Lane 2: Non-transfected lysate.

Storage Buffer

1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction

Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot

Gene Info — IL17RC

Entrez GeneID[84818](#)**GeneBank Accession#**[BC006411.1](#)**Protein Accession#**[AAH06411.1](#)**Gene Name**

IL17RC

Gene Alias

FLJ95963, FLJ96005, IL17-RL, IL17RL, MGC10763

Gene Description

interleukin 17 receptor C

Omim ID[610925](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

This gene encodes a single-pass transmembrane protein that shares limited similarity with the interleukin-17 receptor. Multiple alternatively spliced transcript variants encoding different isoforms have been detected for this gene, but the full-length nature of only three have been determined to date. [provided by RefSeq]

Other Designations

IL17F receptor|interleukin 17 receptor-like