

# NKD1 rabbit monoclonal antibody

Catalog # H00085407-K

Size 100 ug x up to 3

## Specification

Product Description	Rabbit monoclonal antibody raised against a human NKD1 peptide using ARM Technology.
Immunogen	A synthetic peptide of human NKD1 is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
Host	Rabbit
Library Construction	Non-fusion antibody library from rabbit spleen ( <a href="#">ARM Technology</a> ).
Expression	Overexpression vector and transfection into 293H cell line.
Reactivity	Human
Purification	Protein A
Isotype	IgG
Quality Control Testing	Antibody reactive against human NKD1 peptide by ELISA and mammalian transfected lysate by Western Blot.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Deliverable	Up to three rabbit IgG clones of 100 ug each will be delivered to customer.
Note	1. Customer may provide cell or tissue lysate for antibody screening. 2. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering including F(ab) <sub>2</sub> , IgG, scFv and different Fc and non-Fc conjugates per customer request.

## Applications

- Western Blot (Transfected lysate)

[Protocol Download](#)

- ELISA

## Gene Info — NKD1

Entrez GeneID	<a href="#">85407</a>
GeneBank Accession#	<a href="#">NKD1</a>
Gene Name	NKD1
Gene Alias	Naked1
Gene Description	naked cuticle homolog 1 (Drosophila)
Omim ID	<a href="#">607851</a>
Gene Ontology	<a href="#">Hyperlink</a>
Gene Summary	In the mouse, Nkd is a Dishevelled (see DVL1; MIM 601365)-binding protein that functions as a negative regulator of the Wnt (see WNT1; MIM 164820)-beta-catenin (see MIM 116806)-Tcf (see MIM 602272) signaling pathway.[supplied by OMIM]
Other Designations	Dvl-binding protein naked cuticle homolog 1 naked cuticle-1

## Pathway

- [Wnt signaling pathway](#)