


WDR34 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # H00089891-T01


Size 100 uL

Applications



SDS-PAGE Gel

WDR34 transfected lysate.



Western Blot

Lane 1: WDR34 transfected lysate (45.9 KDa)

Lane 2: Non-transfected lysate.

Specification

Transfected Cell Line	293T
Plasmid	pCMV-WDR34 full-length
Host	Human
Theoretical MW (kDa)	45.9
Interspecies Antigen Sequence	Mouse (86)

Quality Control Testing

Transient overexpression cell lysate was tested with Anti-WDR34 antibody ([H00089891-B01](#)) by Western Blots.
SDS-PAGE Gel
WDR34 transfected lysate.
Western Blot
Lane 1: WDR34 transfected lysate (45.9 KDa)
Lane 2: Non-transfected lysate.

Storage Buffer

1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction

Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot

Gene Info — WDR34

Entrez GeneID[89891](#)**GeneBank Accession#**[ENST00000372713](#)**Protein Accession#**[ENSP00000361798](#)**Gene Name**

WDR34

Gene Alias

MGC20486, RP11-216B9.5, bA216B9.3

Gene Description

WD repeat domain 34

Gene Ontology[Hyperlink](#)**Gene Summary**

This gene encodes a member of the WD repeat protein family. WD repeats are minimally conserved regions of approximately 40 amino acids typically bracketed by gly-his and trp-aspartate (GH-WD), which may facilitate formation of heterotrimeric or multiprotein complexes. Members of this family are involved in a variety of cellular processes, including cell cycle progression, signal transduction, apoptosis, and gene regulation. [provided by RefSeq]

Other Designations

OTTHUMP00000022282