


MaxPab@

RNF32 purified MaxPab rabbit polyclonal antibody (D01P)


Catalog # H00140545-D01P Size 100 ug

Applications



Western Blot (Tissue lysate)

RNF32 MaxPab rabbit polyclonal antibody. Western Blot analysis of RNF32 expression in mouse kidney.



Immunoprecipitation

Immunoprecipitation of RNF32 transfected lysate using anti-RNF32 MaxPab rabbit polyclonal antibody and Protein A Magnetic Bead, and immunoblotted with RNF32 purified MaxPab mouse polyclonal antibody (B01P) (H00140545-B01P).

Specification	
Product Description	Rabbit polyclonal antibody raised against a full-length human RNF32 protein.
lmmunogen	RNF32 (AAH28120.1, 1 a.a. ~ 256 a.a) full-length human protein.
Sequence	MLKNKGHSSKKDNLAVNAVALQDHILHDLQLRNLSVADHSKTQVQKKENKSLKRDTKAIIDTGLKK TTQCPKLEDSEKEYVLDPKPPPLTLAQKLGLIGPPPPPLSSDEWEKVKQRSLLQGDSVQPCPICK EEFELRPQVLLSCSHVFHKACLQAFEKFTNKKTCPLCRKNQYQTRVIHDGARLFRIKCVTRIQAYW RGCVVRKWYRNLRKTVPPTDAKLRKNSLKKSSQKSATASCAHTTPTLKSSLQKSISAWP
Host	Rabbit
Reactivity	Human, Mouse



Product Information

Quality Control Testing	Antibody reactive against mammalian tissue lysate.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

Western Blot (Tissue lysate)

RNF32 MaxPab rabbit polyclonal antibody. Western Blot analysis of RNF32 expression in mouse kidney.

Protocol Download

Immunoprecipitation

Immunoprecipitation of RNF32 transfected lysate using anti-RNF32 MaxPab rabbit polyclonal antibody and Protein A Magnetic Bead, and immunoblotted with RNF32 purified MaxPab mouse polyclonal antibody (B01P) (H00140545-B01P).

Protocol Download

Gene Info — RNF32	
Entrez GenelD	<u>140545</u>
GeneBank Accession#	BC028120.1
Protein Accession#	AAH28120.1
Gene Name	RNF32
Gene Alias	FKSG33, HSD15
Gene Description	ring finger protein 32
Omim ID	610241
Gene Ontology	<u>Hyperlink</u>
Gene Summary	The protein encoded by this gene contains two RING ring finger motifs. RING finger motifs are pre sent in a variety of functionally distinct proteins and are known to be involved in protein-DNA or pr otein-protein interactions. This gene was found to be expressed during spermatogenesis, most lik ely in spermatocytes and/or in spermatids. Several alternatively spliced transcript variants exist, b ut their full length natures are not clear. [provided by RefSeq
Other Designations	-