


# RNF32 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # H00140545-T02


Size 100 uL

## Applications



### SDS-PAGE Gel

RNF32 transfected lysate.



### Western Blot

Lane 1: RNF32 transfected lysate ( 28.27 KDa)

Lane 2: Non-transfected lysate.

## Specification

**Transfected Cell Line** 293T

**Plasmid** pCMV-RNF32 full-length

**Host** Human

**Theoretical MW (kDa)** 28.27

**Quality Control Testing** Transient overexpression cell lysate was tested with Anti-RNF32 antibody ([H00140545-B02](#)) by Western Blots.  
 SDS-PAGE Gel  
 RNF32 transfected lysate.  
 Western Blot  
 Lane 1: RNF32 transfected lysate ( 28.27 KDa)  
 Lane 2: Non-transfected lysate.

**Storage Buffer**

1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

**Storage Instruction**

Store at -80°C. Aliquot to avoid repeated freezing and thawing.

## Applications

- Western Blot

## Gene Info — RNF32

**Entrez GeneID**[140545](#)**GeneBank Accession#**[BC028120.1](#)**Protein Accession#**[-](#)**Gene Name**

RNF32

**Gene Alias**

FKSG33, HSD15

**Gene Description**

ring finger protein 32

**Omim ID**[610241](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

The protein encoded by this gene contains two RING ring finger motifs. RING finger motifs are present in a variety of functionally distinct proteins and are known to be involved in protein-DNA or protein-protein interactions. This gene was found to be expressed during spermatogenesis, most likely in spermatocytes and/or in spermatids. Several alternatively spliced transcript variants exist, but their full length natures are not clear. [provided by RefSeq]

**Other Designations**[-](#)